



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : A61K 31/24, 31/42, 31/275, 31/415, 31/495, C07C 211/45, 255/01, C07D 209/18, 231/38, 241/36, 265/34, 471/02, G01N 33/567</p>	<p>A1</p>	<p>(11) International Publication Number: <b>WO 95/21613</b></p> <p>(43) International Publication Date: 17 August 1995 (17.08.95)</p>
<p>(21) International Application Number: PCT/US95/01751</p> <p>(22) International Filing Date: 9 February 1995 (09.02.95)</p> <p>(30) Priority Data: 193,829 9 February 1994 (09.02.94) US</p> <p>(71) Applicants: SUGEN, INC. [US/US]; 515 Galveston Drive, Redwood City, CA 94063 (US). YISSUM RESEARCH DE- VELOPMENT COMPANY OF THE HEBREW UNIVER- SITY IN JERUSALEM [IL/IL]; 46 Jabotinsky Street, P.O. Box 4279, 91042 Jerusalem (IL).</p> <p>(72) Inventors: GAZIT, Aviv; Nof Harim 14, 96190 Jerusalem (IL). LEVITZKI, Alexander; 9617 Fall Bridge Lane, Patomic, MA 02854 (US). APP, Harald; 30 Melrose Court, Hillsborough, CA 94010 (US). TANG, Cho, Peng; 827 Camino Ricardo, Moraga, CA 94556 (US). MCMAHON, Gerald, M.; 1414 Greenwich Street, San Francisco, CA 94109 (US).</p> <p>(74) Agents: CORUZZI, Laura, A. et al.; Pennie &amp; Edmonds, 1155 Avenue of the Americas, New York, NY 10036 (US).</p>	<p>(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KE, KG, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MW, MX, NO, NZ, PL, RO, RU, SD, SI, SK, TJ, TT, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).</p> <p><b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p>(54) Title: COMPOUNDS FOR THE TREATMENT OF DISORDERS RELATED TO VASCULOGENESIS AND/OR ANGIOGENESIS</p>		
<p>(57) Abstract</p> <p>The present invention relates to organic molecules capable of modulating tyrosine kinase signal transduction and particularly KDR/FLK-1 receptor signal transduction in order to regulate and/or modulate vasculogenesis and angiogenesis. The invention is based, in part, on the demonstration that KDR/FLK-1 tyrosine kinase receptor expression is associated with endothelial cells and the identification of vascular endothelial growth factor (VEGF) as the high affinity ligand of FLK-1. These results indicate a major role for KDR/FLK-1 in the signaling system during vasculogenesis and angiogenesis. Engineering of host cells that express FLK-1 and the uses of expressed FLK-1 to evaluate and screen for drugs and analogs of VEGF involved in FLK-1 modulation by either agonist or antagonist activities is also described. The invention also relates to the use of the disclosed compounds in the treatment of disorders, including cancer, diabetes, hemangioma and Kaposi's sarcoma, which are related to vasculogenesis and angiogenesis.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

COMPOUNDS FOR THE  
TREATMENT OF DISORDERS RELATED TO  
VASCULOGENESIS AND/OR ANGIOGENESIS

5        This application is a continuation-in-part and  
divisional of United States Application Serial No.  
08/193,829, filed February 9, 1994, which is a  
continuation-in-part of United States Application  
10       Serial No. 08/038,596, filed March 26, 1993, now  
abandoned, which is a continuation-in-part of United  
States Application Serial No. 07/975,750, filed  
November 13, 1992, now abandoned, all three of which  
are incorporated by reference herein in their  
entirety.

15       1.    INTRODUCTION

          The present invention relates to novel compounds  
capable of modulating and/or regulating tyrosine  
kinase signal transduction. The Applicants have  
20       demonstrated that the murine fetal liver kinase 1  
(FLK-1) receptor and its non-murine counterparts,  
including the human Kinase Insert-Domain-Containing  
Receptor (KDR), play a major role in a tyrosine kinase  
signal transduction system. Polypeptide growth  
25       factors such as vascular endothelial growth factor  
(VEGF) having a high affinity to the KDR/FLK-1  
receptor have been associated with the proliferation  
of endothelial cells and more particularly  
vasculogenesis and/or angiogenesis. Consequently,  
30       compounds affecting the enzymatic function of the  
KDR/FLK-1 receptor may be used to not only regulate,  
modulate and/or inhibit the tyrosine kinase signal  
transduction system, but also the proliferation of  
endothelial cells in processes such as vasculogenesis  
and/or angiogenesis. The present invention is  
35       therefore further directed to the use of compounds

which bind to and/or modulate the activity of the receptors comprising the tyrosine signal transduction system, and more specifically the KDR/FLK-1 receptor, to treat disorders related to vasculogenesis and/or angiogenesis.

## 2. BACKGROUND OF THE INVENTION

**Receptor Tyrosine Kinases.** Receptor tyrosine kinases (RTKs) comprise a large family of transmembrane receptors for polypeptide growth factors with diverse biological activities. The intrinsic function of RTKs is activated upon ligand binding, which results in phosphorylation of the receptor and multiple cellular substrates, and subsequently in a variety of cellular responses. Ullrich & Schlessinger, 1990, *Cell* 61:203-212.

As has been reported by the inventors, RTKs, as well as, more generally, protein tyrosine kinases, play an important role in the control of cell growth and differentiation (for review, see Schlessinger & Ullrich, 1992, *Neuron* 9:383-391). Aberrant expression or mutations in the RTKs have been shown to lead to either uncontrolled cell proliferation (e.g. malignant tumor growth) or to defects in key developmental processes. Consequently, the biomedical community has expended significant resources to discover the specific biological role of members of the RTK family, their function in differentiation processes, their involvement in tumorigenesis and in other diseases, the biochemical mechanisms underlying their signal transduction pathways activated upon ligand stimulation and the development of novel antineoplastic drugs.

At present, at least nineteen (19) distinct RTK subfamilies have been identified. One RTK subfamily is believed to be comprised of the KDR/FLK-1 receptor,

the fetal liver kinase 4 (FLK-4) receptor and the fms-like tyrosine 1 (flt-1) receptor. Each of these receptors was initially believed to be receptors for hematopoietic growth factors.

5       **The KDR/FLK-1 Receptor and VEGF.** Normal vasculogenesis and angiogenesis play important roles in a variety of physiological processes such as embryonic development, wound healing, organ  
10       regeneration and female reproductive processes such as follicle development in the corpus luteum during ovulation and placental growth after pregnancy.

Folkman & Shing, 1992, *J. Biological Chem.* 267(16):10931-34. Uncontrolled vasculogenesis and/or  
15       angiogenesis has been associated with diseases, such as diabetes, as well as malignant solid tumors that rely on vascularization for growth. Klagsburn & Soker, 1993, *Current Biology* 3(10):699-702; Folkham, 1991, *J. Natl., Cancer Inst.* 82:4-6; Weidner, et al., 1991, *New Engl. J. Med.* 324:1-5.

20       Several polypeptides with in vitro endothelial cell growth promoting activity have been identified. Examples include acidic and basic fibroblastic growth factor (FGF), vascular endothelial growth factor  
25       (VEGF) and placental growth factor. Unlike FGF, VEGF has recently been reported to be an endothelial cell specific mitogen. Ferrara & Henzel, 1989, *Biochem. Biophys. Res. Comm.* 161:851-858; Vaisman et al., 1990, *J. Biol. Chem.* 265:19461-19566.

30       Thus, identification of the specific receptors to which VEGF binds is important to understanding of the regulation of endothelial cell proliferation. Two structurally related RTKs have been identified to bind VEGF with high affinity: the flt-1 receptor (Shibuya  
35       et al., 1990, *Oncogene* 5:519-524; De Vries et al., 1992, *Science* 255:989-991) and the KDR/FLK-1 receptor,

discussed herein. Consequently, it had been surmised that RTKs may have a role in the modulation and regulation of endothelial cell proliferation.

5 As has only been recently contemplated, evidence, such as information set forth in U.S. Application Serial Nos. 08/193,829, 08/038,596 and 07/975,750, strongly suggest that VEGF is not only responsible for endothelial cell proliferation, but also is the prime  
10 regulator of normal and pathological angiogenesis. See generally, Klagsburn & Soker, 1993, *Current Biology* 3(10):699-702; Houck, et al., 1992, *J. Biol. Chem.* 267:26031-26037.

**Identification Of Agonists And Antagonists To The KDR/FLK-1 Receptor.** In view of the surmised  
15 importance of RTKs to the control, regulation and modulation of endothelial cell proliferation and potentially vasculogenesis and/or angiogenesis, many attempts have been made to identify RTK "inhibitors" using a variety of approaches, including the use of  
20 mutant ligands (U.S. Application No. 4,966,849), soluble receptors and antibodies (Application No. WO 94/10202; Kendall & Thomas, 1994, *Proc. Nat'l Acad. Sci.* 90:10705-09; Kim, et al., 1993, *Nature* 362:841-844), RNA ligands (Jellinek, et al., 19\_\_,  
25 *Biochemistry* 33:10450-56), protein kinase C inhibitors (Schuchter, et al., 1991, *Cancer Res.* 51:682-687); Takano, et al., 1993, *Mol. Bio. Cell* 4:358A; Kinsella, et al., 1992, *Exp. Cell Res.* 199:56-62; Wright, et al., 1992, *J. Cellular Phys.* 152:448-57) and tyrosine  
30 kinase inhibitors (WO 94/03427; WO 92/21660; WO 91/15495; WO 94/14808; U.S. Patent No. 5,330,992; Mariani, et al., 1994, *Proc. Am. Assoc. Cancer Res.* 35:2268).

35 More recently, attempts have been made to identify small molecules which act as tyrosine kinase


inhibitors. For example, bis monocyclic, bicyclic or heterocyclic aryl compounds (PCT WO 92/20642), vinylene-aza-indole derivatives (PCT WO 94/14808) and 1-cyclopropyl-4-pyridyl-quinolones (U.S. Patent No. 5,330,992) have been described generally as tyrosine kinase inhibitors. Styryl compounds (U.S. Patent No. 5,217,999), styryl-substituted pyridyl compounds (U.S. Patent No. 5,302,606), certain quinazoline derivatives (EP Application No. 0 566 266 A1), seleoindoles and selenides (PCT WO 94/03427), tricyclic polyhydroxylic compounds (PCT WO 92/21660) and benzylphosphonic acid compounds (PCT WO 91/15495) have been described as compounds for use as tyrosine kinase inhibitors for use in the treatment of cancer. None of these compounds, however, have been previously associated with the enzymatic function of the KDR/FLK-1 receptor. Likewise, none of these compounds have been associated with regulation of vasculogenesis and/or angiogenesis.

The identification of effective small compounds which specifically inhibit tyrosine signal transduction by modulating the activity of RTKs and particularly the KDR/FLK-1 receptor to regulate and modulate vasculogenesis and/or angiogenesis is therefore desirable and the object of this invention.

### 3. SUMMARY OF THE INVENTION

The present invention relates to organic molecules capable of modulating tyrosine signal transduction and the use of such molecules to inhibit or promote angiogenesis and/or vasculogenesis. Generally, the compounds of the instant invention are derivatives of quinazoline, quinoxiline, substituted aniline, isoxazoles, acrylonitrile and phenylacrylonitrile compounds. More specifically, the

invention is generally directed to compounds having the formulae:



10 and pharmaceutically acceptable salts thereof, wherein:

- R<sub>1</sub> is isopropyl, t-butyl, I, Br, OH or methyl,  
 R<sub>2</sub> is OH,  
 15 R<sub>3</sub> is 2-propyl, t-butyl, OH, H or methyl, and  
 R<sub>4</sub> is (1-phenyl)-n-propylaminocarbonyl, (E) 1-cyano-2-[(3,5-diisopropyl-4-hydroxy)phenyl]ethenyl-sulfonyl, aminothiocabonyl, cyanomethylsulfonyl, (3-amino-4-cyano)pyrazo-5-yl, phenyl-n-  
 20 propylaminocarbonyl, (E) 1-cyano-2-[(5-bromo-3,4-dihydroxy)phenyl]ethenylsulfonyl, (1-phenyl)-n-propylaminothiocarbonyl, cyano, (E) [[[4-[1-cyano-2(3,4-dihydroxy)phenyl]ethenyl]carbonylamino]-n-butyl]aminocarbonyl, benzylaminocarbonyl, 2[[2-cyano-1-(3,4-dihydroxy)phenyl]ethylenyl]]sulfonyl, [(3,4-  
 25 dihydroxy)phenyl]carbonyl, (E) [[[4-[1-cyano-2(3,4-dihydroxy)phenyl]ethenyl]carbonylamino]-ethyl]aminocarbonyl or hydroxycarbonyl;


-OR-

30

35



B.



and pharmaceutically acceptable salts thereof,  
wherein:

R<sub>1</sub> is CH<sub>3</sub> or H,

R<sub>2</sub> is CH<sub>3</sub> or H, or alternatively,


R<sub>1</sub> and R<sub>2</sub> form a phenyl ring (CHCHCHCH),

R<sub>3</sub> is H or formyl or chloro, and

R<sub>4</sub> is phenyl, (3,4-dihydroxy)phenyl, (4-iodophenyl)amino, (3,4-dichlorophenyl)amino, (3-chlorophenyl)amino, (4-bromophenyl)amino or n-propylamino;

-OR-

C.



and pharmaceutically acceptable salts thereof,  
wherein:

R<sub>1</sub> is OCH<sub>3</sub>, CH<sub>3</sub> or H,


R<sub>2</sub> is OCH<sub>3</sub> or H,

R<sub>3</sub> is H or chloro, and

$R_4$  is (3-chlorophenyl)amino, (4-methylphenyl)mecapto, (4-iodophenyl)amino or (3-hydroxyphenyl)amino;

-OR-

D.



and pharmaceutically acceptable salts thereof, wherein:


$R_1$  is 4-hydroxy-phenyl

$R_2$  is benzyl, and

$R_3$  is  $\text{CH}_3$ ;

-OR-

E.




and pharmaceutically acceptable salts thereof, wherein:

$R_1$  is (3-trifluormethyl)phenyl and

$R_2$  is (2-chlorophenyl)aminothiocarbonyl;

-OR-

F.



and pharmaceutically acceptable salts thereof,  
wherein:


$R_1$  is O,

$R_2$  is (3,4-dihydroxyphenyl), and

$R_3$ ,  $R_4$  and  $R_5$  are H;

-OR-

G.



and pharmaceutically acceptable salts thereof,  
wherein:


$R_1$  is 4-(1-nitro)thiophene or indol-3-yl or indol-5-yl and

$R_2$  is aminothiocarbonyl, (3-amino-4-cyano)pyrazol-5-yl or (3,4-dihydroxyphenyl)carbonyl;

-OR-

H.

5



and pharmaceutically acceptable salts thereof,  
 wherein:

10

$R_1$  is 2,5-dihydroxylbenzyl,


$R_2$  is H,

$R_3$  is methoxycarbonyl, and

$R_4$  is H;

15

-OR-



I.

20

and pharmaceutically acceptable salts thereof,  
 wherein:

25

$R_1$  is H,


$R_2$  is (4-trifluoromethyl)phenyl, and

$R_3$  is methyl;

-OR-

30

J.



and pharmaceutically acceptable salts thereof, wherein  
 $R_1$  is (3-chloro)phenylamino.

35

The present invention is further directed to pharmaceutical compositions comprising a pharmaceutically effective amount of the above-described compounds and a pharmaceutically acceptable carrier or excipient. Such a composition is believed to affect the enzymatic activity of the KDR/FLK-1 receptor by, *inter alia*, inhibiting the signal transduced by interaction between KDR/FLK-1 and vascular endothelial growth factor (VEGF) which may be useful in inhibition of diseases related to vasculogenesis and/or angiogenesis, including diabetes and cancer. Alternatively, such composition may act directly on the cells responsible for the disease (e.g. tumor cells). More particularly, the compositions of the present invention may be included in methods for treating, among other diseases, diabetic retinopathy, glioma, melanoma, Kaposi's sarcoma, hemangioma and ovarian, breast, lung, pancreatic, prostate, colon and epidermoid cancer. Such a composition, if used to modulate, rather than inhibit, vasculogenesis and/or angiogenesis may also be useful in the promotion of wound healing.

Finally, the present invention is also directed to methods for treating diseases related to pathological vasculogenesis and/or angiogenesis, including but not limited to diabetes, diabetic retinopathy, rheumatoid arthritis, hemangioma and cancer and more particularly cancer related to solid cell tumor growth (e.g., glioblastoma, melanoma and Kaposi's sarcoma and ovarian, lung, mammary, prostate, pancreatic, colon and epidermoid carcinoma).

### 3.1. Definitions

"Pharmaceutically acceptable acid addition salt" refers to those salts which retain the biological effectiveness and properties of the free bases and which are obtained by reaction with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid and the like.

"Alkyl" refers to saturated or unsaturated branched or straight chain hydrocarbon radical. Typical alkyl groups includes methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tertiary butyl, pentyl, hexyl and the like.

### 4. DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to compounds capable of regulating and/or modulating tyrosine signal transduction and more particularly KDR/FLK-1 receptor signal transduction in order to inhibit or promote angiogenesis and/or vasculogenesis.

The present invention is based, in part, on the discovery that the KDR/FLK-1 tyrosine kinase receptor, and RTKs more generally, is expressed on the surface of endothelial cells and may play a role in endothelial cell growth, including solid cell tumor growth. The invention is also based on the identification of VEGF as a high affinity ligand of KDR/FLK-1 and the characterization of KDR/FLK-1 as an RTK rather than a hematopoietic receptor. Thus, the surmised role of VEGF in endothelial cell proliferation and migration during angiogenesis and vasculogenesis indicate an important role for the KDR/FLK-1 in these processes.

The invention is further based on the observation that diseases such as diabetes mellitus (Folkman, 1987, in XIth Congress of Thrombosis and Haemostasis (Verstraeta, et al., eds.) pp. 583-596, Leuven University Press, Leuven) and arthritis, as well as malignant tumor growth may result from uncontrolled angiogenesis. See e.g., Folkman, 1971, N. Engl. J. Med. 285:1182-1186. Finally, the invention is based upon the discovery and design of compounds that inhibit, prevent, or interfere with the signal transduced by KDR/FLK-1 when activated by ligands such as VEGF. Although it is therefore believed that the compounds of the present invention act on a receptor or oother component along the tyrosine kinase signal transduction pathway, the compounds may also act directly on the tumors cells that result from uncontrolled angiogenesis.

For purposes of this application, although the nomenclature of the human and murine counterparts of the generic "flk-1" receptor differ, they are, in many respects, interchangeable. The murine receptor, FLK-1, and its human counterpart, KDR, share a sequence homology of 93.4% within the intracellular domain. Likewise, murine FLK-1 binds human VEGF with the same affinity as mouse VEGF, and accordingly, is activated by the ligand derived from either species. Millauer et al., 1993, *Cell* 72:835-846; Quinn et al., 1993, *Proc. Natl. Acad. Sci. USA* 90:7533-7537. FLK-1 also associates with and subsequently tyrosine phosphorylates human RTK substrates (e.g., PLC- $\gamma$  or p85) when coexpressed in 293 cells (human embryonal kidney fibroblasts).

Models which rely upon the FLK-1 receptor therefore are directly applicable to understanding the KDR receptor. For example, use of the murine FLK-1

receptor in methods to identify compounds which regulate the signal transduction pathway are directly applicable to the identification of compounds which may be used to regulate the human signal transduction pathway, and more specifically, activity related to the KDR receptor. Angiogenesis is a very complex process involving the invasion of endothelial cells into the nonvascularized tissue. No *in vitro* model exists which mimics exactly this multistep process comprising the degradation of the basal membrane surrounding the endothelial cells, migration into the perivascular stroma and eventually proliferation and formation of the new vascular sprout. Thus, chemical compounds identified as inhibitors of KDR/FLK-1 *in vitro*, will be confirmed in suitable *in vivo* models. Both *in vivo* mouse and rat animal models have been demonstrated to be of excellent value for the examination of the clinical potential of agents acting on the KDR/FLK-1 induced signal transduction pathway.

In sum, the receptors to which VEGF specifically binds are an important and powerful therapeutical target for the regulation and modulation of vasculogenesis and/or angiogenesis and a variety of severe diseases which involve abnormal cellular growth caused by such processes. Plowman, et al., 1994, *DN&P* 7(6):334-339. More particularly, the KDR/FLK-1 receptor's high specificity and role in the neovascularization make it a very distinct and powerful target for therapeutic approaches for the treat cancer and other diseases which involve the uncontrolled formation of blood vessels.

This invention is therefore directed to compounds which regulate, modulate and/or inhibit vasculogenesis and/or angiogenesis by affecting the enzymatic activity of the KDR/FLK-1 receptor and interfering



with the signal transduced by KDR/FLK-1. More particularly, the present invention is directed to compounds which regulate, modulate and/or inhibit the KDR/FLK-1 mediated signal transduction pathway as a therapeutic approach to cure many kinds of solid tumors, including but not limited to glioblastoma, melanoma and Kaposi's sarcoma, and ovarian, lung, mammary, prostate, pancreatic, colon and epidermoid carcinoma. In addition, data suggest the administration of compounds which inhibit the KDR/FLK-1 mediated signal transduction pathway to the treatment of hemangioma and diabetic retinopathy.

The invention also relates to the inhibition of vasculogenesis and angiogenesis via other receptor-mediated pathways, including the pathway comprising the highly related flt-1 receptor. Receptor tyrosine kinase mediated signal transduction is initiated by extracellular interaction with a specific growth factor (ligand), followed by receptor dimerization, transient stimulation of the intrinsic protein tyrosine kinase activity and autophosphorylation. Binding sites are thereby created for intracellular signal transduction molecules and lead to the formation of complexes with a spectrum of cytoplasmic signalling molecules that facilitate the appropriate cellular response. (E.g., cell division, metabolic effects to the extracellular microenvironment). See, Schlessinger and Ullrich, 1992, *Neuron* 9:1-20.

It has been shown that tyrosine autophosphorylation sites in growth factor receptors, such as KFR/FLK-1 and flt-1, function as high-affinity binding sites for SH2 (*src* homology) domains of signaling molecules. Fantl et al., 1992, *Cell* 69:413-423; Songyang et al., 1994, *Mol. Cell. Biol.* 14:2777-2785; Songyang et al., 1993, *Cell* 72:767-778; and

Koch et al., 1991, *Science* 252:668-678. Several intracellular substrate proteins that associate with receptor tyrosine kinases have been identified. They may be divided into two principal groups: (1) substrates which have a catalytic domain; and (2) substrates which lack such domain but serve as adapters and associate with catalytically active molecules. Songyang et al., 1993, *Cell* 72:767-778. The specificity of the interactions between receptors, such as KDR/FLK-1 and flt-1, and SH2 domains of their substrates is determined by the amino acid residues immediately surrounding the phosphorylated tyrosine residue. Differences in the binding affinities between SH2 domains and the amino acid sequences surrounding the phosphotyrosine residues on particular receptors are consistent with the observed differences in their substrate phosphorylation profiles. Songyang et al., 1993, *Cell* 72:767-778. These observations suggest that the function of each receptor tyrosine kinase is determined not only by its pattern of expression and ligand availability but also by the array of downstream signal transduction pathways that are activated by a particular receptor. Thus, autophosphorylation provides an important regulatory step which determines the selectivity of signaling pathways recruited by specific growth factor receptors, as well as differentiation factor receptors.

The close homology of the intracellular regions of KDR/FLK-1 with that of the PDGF- $\beta$ -Receptor (50.3% homolgy) and/or the highly related flt-1 receptor indicates the induction of overlapping signal transduction pathways. For example, for the PDGF- $\beta$ -Receptor, members of the src family (Twamley et al., 1993, *Proc. Natl. Acad. Sci. USA* 90:7696-7700),


phosphatidylinositol-3'-kinase (Hu et al., 1992, *Mol. Cell. Biol.* 12:981-990), phospholipase C- $\gamma$  (Kashishian & Cooper, 1993, *Mol. Cell. Biol.* 4:49-51), ras-GTPase-activating protein, (Kashishian et al., 1992, *EMBO J.* 11:1373-1382), PTP-1D/syp (Kazlauskas et al., 1993, *Proc. Natl. Acad. Sci. USA* 90:6939-6943), Grb2 (Arvidsson et al., 1994, *Mol. Cell. Biol.* 14:6715-6726), and the adapter molecules Shc and Nck (Nishimura et al., 1993, *Mol. Cell. Biol.* 13:6889-6896), have been shown to bind to regions involving different autophosphorylation sites. See generally, review, please see Claesson-Welsh, 1994, *Prog. Growth Factor Res.* 5:37-54. Thus, it is likely that signal transduction pathways activated by KDR/FLK-1 include the ras pathway (Rozakis et al., 1992, *Nature* 360:689-692), the PI-3'-kinase pathway and the src-mediated and plcy-mediated pathways. Each of these pathways may play a critical role in the angiogenic and/or vasculogenic effect of KDR/FLK-1 in endothelial cells. Consequently, the present invention is also directed to the use of the organic compounds discussed herein to modulate angiogenesis and vasculogenesis as such processes are controlled by these pathways.

#### 4.1. The Compounds

The invention is generally directed to compounds and/or compositions comprising compounds having the formulae:

A.

5



10 and pharmaceutically acceptable salts thereof,  
wherein:

$R_1$  is isopropyl, t-butyl, I, Br, OH or methyl,

$R_2$  is OH,

$R_3$  is 2-propyl, t-butyl, OH, H or methyl, and

15

$R_4$  is (1-phenyl)-n-propylaminocarbonyl, (E) 1-cyano-2-[(3,5-diisopropyl-4-hydroxy)phenyl]ethenyl-sulfonyl, aminothiocabonyl, cyanomethylsulfonyl, (3-amino-4-cyano)pyrazo-5-yl, phenyl-n-

20


propylaminocarbonyl, (E) 1-cyano-2-[(5-bromo-3,4-dihydroxy)phenyl]ethenylsulfonyl, (1-phenyl)-n-propylaminothiocarbonyl, cyano, (E) [[[4-[1-cyano-2(3,4-dihydroxy)phenyl]ethenyl]carbonylamino]-n-butyl]aminocarbonyl, benzylaminocarbonyl, 2[[2-cyano-1-(3,4-dihydroxy)phenyl]ethylenyl]]sulfonyl, [(3,4-dihydroxy)phenyl]carbonyl, (E) [[[4-[1-cyano-2(3,4-dihydroxy)phenyl]ethenyl]carbonylamino]-ethyllaminocarbonyl or hydroxycarbonyl;

25

-OR-

30

B.



35

and pharmaceutically acceptable salts thereof,  
wherein:

$R_1$  is  $\text{CH}_3$  or H,

$R_2$  is  $\text{CH}_3$  or H, or alternatively,


$R_1$  and  $R_2$  form a phenyl ring ( $\text{CHCHCHCH}$ ),

$R_3$  is H or formyl or chloro, and

$R_4$  is phenyl, (3,4-dihydroxy)phenyl, (4-iodophenyl)amino, (3,4-dichlorophenyl)amino, (3-chlorophenyl)amino, (4-bromophenyl)amino or n-propylamino;

-OR-

C.



and pharmaceutically acceptable salts thereof,  
wherein:

$R_1$  is  $\text{OCH}_3$ ,  $\text{CH}_3$  or H,

$R_2$  is  $\text{OCH}_3$  or H,


$R_3$  is H or chloro, and

$R_4$  is (3-chlorophenyl)amino, (4-methylphenyl)mecapto, (4-iodophenyl)amino or (3-hydroxyphenyl)amino;

-OR-

D.

and  
pharmaceutic  
ally  
5 acceptable  
salts  
thereof,  
wherein:




10 R<sub>1</sub> is 4-hydroxy-phenyl

R<sub>2</sub> is benzyl, and

R<sub>3</sub> is CH<sub>3</sub>;

E.




20 and pharmaceutically acceptable salts thereof,  
wherein:

R<sub>1</sub> is (3-trifluormethyl)phenyl and

R<sub>2</sub> is (2-chlorophenyl)aminothiocarbonyl;

F.



35 and pharmaceutically acceptable salts thereof,  
wherein:

$R_1$  is O,


$R_2$  is (3,4-dihydroxy)phenyl, and

$R_3$ ,  $R_4$  and  $R_5$  are H;

5

G.

10



and pharmaceutically acceptable salts thereof,  
wherein:


15

$R_1$  is 4-(1-nitro)thiophene or indol-3-yl or indol-5-yl and

$R_2$  is aminothiocarbonyl, (3-amino-4-cyano)pyrazol-5-yl or (3,4-dihydroxyphenyl)carbonyl;

20

H.



25

and pharmaceutically acceptable salts thereof,  
wherein:

30

$R_1$  is 2,5-dihydroxybenzyl,

$R_2$  is H,

$R_3$  is methoxycarbonyl, and


$R_4$  is H;

35

I.

-OR-


5 and  
pharmaceutic  
ally  
acceptable  
salts thereof, wherein:



10 R<sub>1</sub> is H,  
R<sub>2</sub> is (4-trifluoromethyl)phenyl, and  
R<sub>3</sub> is methyl;

-OR-

15 J.



20 and pharmaceutically acceptable salts thereof, wherein  
R<sub>1</sub> is (3-chloro)phenylamino.

25 The chemical formulae referred herein may exhibit  
the phenomenon of tautomerism. As the formulae  
drawings within this specification can only represent  
one of the possible tautomeric forms, it should be  
understood that the invention encompasses any  
tautomeric form which possesses the ability to  
regulate and/or modulate vasculogenesis and/or  
angiogenesis and is not limited to any one tautomeric  
30 form utilized within the formulae drawings.

In addition to the above compounds and their  
pharmaceutically acceptable salts, the invention is  
further directed, where applicable, to solvated as  
well as unsolvated forms of the compounds (e.g.  
35



hydrated forms) having the ability to regulate and/or modulate vasculogenesis and/or angiogenesis.

The compounds described above may be prepared by any process known to be applicable to the preparation of chemically-related compounds. Suitable processes are illustrated by the following representative examples. Necessary starting materials may be obtained by standard procedures of organic chemistry.

#### 4.2. Pharmaceutical Formulations And Routes Of Administration

The identified compounds can be administered to a human patient, by itself, or in pharmaceutical compositions where it is mixed with suitable carriers or excipient(s) at doses to treat or ameliorate a variety of disorders, including solid cell tumor growth, including Kaposi's sarcoma, glioblastoma, and melanoma and ovarian, lung, mammary, prostate, pancreatic, colon and epidermoid carcinoma, diabetes, diabetic retinopathy, hemangioma and rheumatoid arthritis. A therapeutically effective dose further refers to that amount of the compound sufficient to result in amelioration of symptoms of uncontrolled vasculogenesis and angiogenesis. Techniques for formulation and administration of the compounds of the instant application may be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, PA, latest edition.

##### 4.2.1. Routes Of Administration.

Suitable routes of administration may, for example, include oral, rectal, transmucosal, or intestinal administration; parenteral delivery, including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections.

Alternately, one may administer the compound in a local rather than systemic manner, for example, via injection of the compound directly into a solid tumor, often in a depot or sustained release formulation.

Furthermore, one may administer the drug in a targeted drug delivery system, for example, in a liposome coated with tumor-specific antibody. The liposomes will be targeted to and taken up selectively by the tumor.

#### 4.2.2. Composition/Formulation.

The pharmaceutical compositions of the present invention may be manufactured in a manner that is itself known, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

Pharmaceutical compositions for use in accordance with the present invention thus may be formulated in conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen.

For injection, the agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological saline buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

For oral administration, the compounds can be formulated readily by combining the active compounds with pharmaceutically acceptable carriers well known

in the art. Such carriers enable the compounds of the invention to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated.

5       Pharmaceutical preparations for oral use can be obtained solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if  
10       desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize  
15       starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium  
20       carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium  
25       alginate.

      Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene  
25       glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize  
30       different combinations of active compound doses.

      Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The  
35       push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such

as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils,  
5 liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. All formulations for oral administration should be in dosages suitable for such administration.

10 For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray  
15 presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol  
20 the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of e.g. gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or  
25 starch.

The compounds may be formulated for parenteral administration by injection, e.g., by bolus injection or continuous infusion. Formulations for injection  
30 may be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as  
35 suspending, stabilizing and/or dispersing agents.

Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

A pharmaceutical carrier for the hydrophobic compounds of the invention is a cosolvent system comprising benzyl alcohol, a nonpolar surfactant, a

water-miscible organic polymer, and an aqueous phase. The cosolvent system may be the VPD co-solvent system. VPD is a solution of 3% w/v benzyl alcohol, 8% w/v of the nonpolar surfactant polysorbate 80, and 65% w/v polyethylene glycol 300, made up to volume in absolute ethanol. The VPD co-solvent system (VPD:5W) consists of VPD diluted 1:1 with a 5% dextrose in water solution. This co-solvent system dissolves hydrophobic compounds well, and itself produces low toxicity upon systemic administration. Naturally, the proportions of a co-solvent system may be varied considerably without destroying its solubility and toxicity characteristics. Furthermore, the identity of the co-solvent components may be varied: for example, other low-toxicity nonpolar surfactants may be used instead of polysorbate 80; the fraction size of polyethylene glycol may be varied; other biocompatible polymers may replace polyethylene glycol, e.g. polyvinyl pyrrolidone; and other sugars or polysaccharides may substitute for dextrose.

Alternatively, other delivery systems for hydrophobic pharmaceutical compounds may be employed. Liposomes and emulsions are well known examples of delivery vehicles or carriers for hydrophobic drugs. Certain organic solvents such as dimethylsulfoxide also may be employed, although usually at the cost of greater toxicity. Additionally, the compounds may be delivered using a sustained-release system, such as semipermeable matrices of solid hydrophobic polymers containing the therapeutic agent. Various of sustained-release materials have been established and are well known by those skilled in the art. Sustained-release capsules may, depending on their chemical nature, release the compounds for a few weeks up to over 100 days. Depending on the chemical nature

and the biological stability of the therapeutic reagent, additional strategies for protein stabilization may be employed.

5       The pharmaceutical compositions also may comprise suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients include but are not limited to calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers such as  
10 polyethylene glycols.

      Many of the KDR/FLK-1 receptor modulating compounds of the invention may be provided as salts with pharmaceutically compatible counterions.  
15 Pharmaceutically compatible salts may be formed with many acids, including but not limited to hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents that are the corresponding  
20 free base forms.

#### 4.2.3.       Effective Dosage.

      Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in an  
25 effective amount to achieve its intended purpose. More specifically, a therapeutically effective amount means an amount effective to prevent development of or to alleviate the existing symptoms of the subject being treated. Determination of the effective amounts  
30 is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein.

      For any compound used in the method of the invention, the therapeutically effective dose can be  
35 estimated initially from cell culture assays. For example, a dose can be formulated in animal models to

achieve a circulating concentration range that includes the IC50 as determined in cell culture (i.e., the concentration of the test compound which achieves a half-maximal inhibition of the RTK activity). Such information can be used to more accurately determine useful doses in humans.

A therapeutically effective dose refers to that amount of the compound that results in amelioration of symptoms or a prolongation of survival in a patient. Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio between LD50 and ED50. Compounds which exhibit high therapeutic indices are preferred. The data obtained from these cell culture assays and animal studies can be used in formulating a range of dosage for use in human. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. (See e.g. Fingl et al., 1975, in "The Pharmacological Basis of Therapeutics", Ch. 1 p1).

Dosage amount and interval may be adjusted individually to provide plasma levels of the active moiety which are sufficient to maintain the FLK-1 receptor-inhibitory effects. Usual patient dosages



for systemic administration range from 1 - 2000 mg/day, commonly from 1 - 250 mg/day, and typically from 10 - 150 mg/day. Stated in terms of patient body weight, usual dosages range from 0.02 - 25 mg/kg/day, commonly from 0.02 - 3 mg/kg/day, typically from 0.2 - 1.5 mg/kg/day. Stated in terms of patient body surface areas, usual dosages range from 0.5 - 1200 mg/m<sup>2</sup>/day, commonly from 0.5 - 150 mg/m<sup>2</sup>/day, typically from 5 - 100 mg/m<sup>2</sup>/day.

Dosage amount and interval may be adjusted individually to provide plasma levels of the active moiety which are sufficient to maintain the FLK-1 receptor-inhibitory effects. Usual average plasma levels should be maintained within 50 - 5000 µg/ml, commonly 50 - 1000 µg/ml, and typically 100 - 500 µg/ml

In cases of local administration or selective uptake, the effective local concentration of the drug may not be related to plasma concentration.

The amount of composition administered will, of course, be dependent on the subject being treated, on the subject's weight, the severity of the affliction, the manner of administration and the judgment of the prescribing physician.

#### 4.2.4. Packaging

The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack may for example comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration. Compositions comprising a compound of the invention formulated in a compatible pharmaceutical carrier may also be prepared, placed in an appropriate container, and

labelled for treatment of an indicated condition. Suitable conditions indicated on the label may include treatment of a tumor, such as a glioma or glioblastoma and inhibition of angiogenesis.

5

**5. EXAMPLE: Compound Synthesis**

**5.1. Synthesis Of (E)-2-aminothiocarbonyl-3-(3,5-di-t-butyl-4-hydroxyphenyl)acrylonitrile**

A preferred method of synthesis of (E)-2-aminothiocarbonyl-3 (3,5-di-t-butyl-4-hydroxyphenyl)acrylonitrile (Compound 1) is as follows: The compound was prepared as generally described by Ohmichi et al., 1993, *Biochemistry* 32:4650. A mixture of 0.47 g of 3,5-Di-tert-butyl-4-hydroxybenzaldehyde, 0.2 g of thiocyanacetamide and 30 mg of  $\beta$ -alanine in 40 ml of ethanol was refluxed for 6 hours. Water and HCl were added, and the reaction mixture was extracted with ethyl acetate. Evaporation gave 0.34 g (54% yield) of a yellow solid having a melting point of 210°C.

20

The product gave the following analytical data: NMR (acetone- $d_6$ )  $\delta$  8.47 (1H, s, vinyl), 8.02 (2 H, s), 1.48 (18 H, s); MS m/e 316 (M<sup>+</sup>, 100), 303 (35), 301 (99), 268 (16), 260 (M (CH<sub>3</sub>)<sub>2</sub>=C, 45), 245 (17), 228 (22), 219 (52), 203 (10), 143 (11), 129 (11).

25

**5.2. Synthesis Of (E)-2-cyano-3-(3-iodo-4,5-hydroxyphenyl)acrylonitrile**

A preferred method of synthesis of (E)-1-cyano-3-(3-iodo-4,5-hydroxyphenyl)acrylonitrile (Compound 2) is as follows: The compound was prepared as described in Ohmichi et al., 1993, *Biochemistry* 32:4650 in two steps. First, 3-methoxy-4-hydroxy-5-iodobenzylidene malononitrile was prepared by adding 3 drops of piperidine to 1.4 g of 5-iodovanillin and 0.4 g of malononitrile in 25 ml of ethanol and refluxing

35

the mixture for 4 hours. 0.8 g (49% yield) of a yellow solid resulted.

The 3-methoxy-4-hydroxy-5-

5 (iodobenzylidene)malononitrile product gave the following analytical data: mp 188°C; NMR (CDCl<sub>3</sub>)  $\delta$  7.76 (1 H, J=1.8 Hz, H<sub>6</sub>), 7.65 (1 H, d, J=1.8 Hz, H<sub>2</sub>), 7.56 (1 H, s, vinyl), 6.85 (1 H, s, OH), 3.99 (3 H, s, OCH<sub>3</sub>); MS m/e 327 (13), 326 (M<sup>+</sup>, 100), 283 (18), 128 (35), 101 (22).

10

Next, (3-methoxy-4-hydroxy-5-iodobenzylidene)malononitrile (0.65 g) and 0.6 ml of boron tribromide (BBr<sub>3</sub>) in 40 ml of dichloromethane were stirred under argon for 1 hour at room temperature. Water was added, and the reaction mixture was extracted with ethyl acetate to give 0.46 g (73% yield) of a light-red solid (yellow in solution) having a melting point of 105°C.

15

The final product gave the following analytical data: NMR (acetone-d<sub>6</sub>)  $\delta$  8.03 (1 H, s, vinyl), 7.88 (1 H, d, J=2.3 Hz, H<sub>2</sub>), 7.72 (1 H, d, J=2.3 Hz, H<sub>6</sub>); MS m/e 312 (M<sup>+</sup>, 38), 254 (74), 185 (M-I, 27), 158 (M-I-HCN, 11), 157 (64), 130 (19), 129 (23), 127 (100).

20

### 25 5.3. Synthesis Of (E)-2-(3-phenyl-n-propylaminocarbonyl)-3-(3-bromo-4,5-dihydroxyphenyl)acrylonitrile

A preferred method of synthesis of (E)-2-(3-phenyl-n-propylaminocarbonyl)-3-(3-bromo-4,5-dihydroxyphenyl)acrylonitrile (Compound 3) is as follows:

30 1. A mixture of 0.69 g of 2.5 mM 5-iodovanillin, 0.5 g of N-3-phenyl-n-propyl cyanoacetamide and 50 mg  $\beta$ -alanine in 30 ml ethanol was refluxed for 5 hours. Evaporation gave an oil which was triturated with benzene-hexane and filtered to give 0.82 g of a bright yellow solid (71% yield)

35

having a melting point of 83°C. Notably, the compound was observed to partially deteriorate over time when stored at room light. It is therefore preferred that the compound be stored as a solid and protected from light. The product gave the following analytical data: NMR (CDCl<sub>3</sub>) δ 8.12 (1H, s), 7.75 (1H, d, J=2.0 Hz), 7.68 (1H, d, J=2.0 Hz), 7.30-7.10 (5H, m), 3.96 (3H, s, OCH<sub>3</sub>), 3.45 (2H, q, J=6.0 Hz), 2.70 (2H, t, J=6.0 Hz), 1.95 (2H, quin, J=6.0 Hz). MS m/e 462 (M<sup>+</sup>, 53), 357 (M-CH<sub>2</sub>)<sub>3</sub>Ph, 18), 335 (M-I, 100), 327 (M-NH(CH<sub>2</sub>)<sub>3</sub> Ph, 31).

2. 0.5 g of the compound of step 1 (3-methoxy-4-hydroxy-5-iodo α-cis cinnamone (3'-phenylpropane) amide) and 0.4 ml of BBr<sub>3</sub> in 30 ml dichloromethane were then stirred at room temperature for 1.5 hours. Water was added and the reaction extracted with ethyl acetate. Evaporation and trituration with benzene-hexane gave 0.3 g of a light brown solid (63% yield) having a melting temperature of 184°C.

The product gave the following analytical data: NMR (acetone d<sub>6</sub>) δ 8.01 (1H, s vinyl), 7.88 (1H, d, J=2.0 Hz), 7.66 (1H, d, J=2.0 Hz), 7.30 (5H, m, Ph), 3.42 (2H, t, J=6.0 Hz), 2.70 (2H, t, J=6.0 Hz), 1.96 (2H, quin., J=6.0 Hz). MS m/e 448 (M<sup>+</sup>, 3%), 321 (M-I, 8), 217 (21), 201 (33), 118 (100), m/e.

#### 5.4. Synthesis Of (E)-2-[(3-amino-4-cyano)pyrazo-5-yl]-3-(3,5-di-t-butyl-4-hydroxyphenyl)acrylonitrile

A preferred method of synthesis of (E)-2-[(3-amino-4-cyano)pyrazo-5-yl]-3-(3,5-di-t-butyl-4-hydroxyphenyl)acrylonitrile (Compound 4) is as follows: A mixture of 0.7 g of 3,5-di-t-butyl-4-hydroxybenzaldehyde, 0.46 g of 3-amino-4-cyano-5-cyanomethyl pyrazole (prepared according to Carboni et

al., 1958, *J. Chem. Soc.*, 80:2838) and 40 mg of  $\beta$ -alanine was refluxed for 15 hours. Cooling and filtering gave 0.5 g (46% yield) of yellow solid having a melting point of 255°C.

The product gave the following analytical data:  
NMR (CDCl<sub>3</sub>)  $\delta$  7.92 (1H, S, vinyl), 7.80 (2H, S), 5.76 (1H, S, OH), 3.75 (2H, Br, S, NH<sub>2</sub>), 1.48 (18H, S). MS m/e 364 (M+1, 28), 363 (M+, 100%), 348 (M-CH<sub>3</sub>, 58%), 292 (M-56-CH<sub>3</sub>, 31%), 147 (41%), m/e.

#### 5.5. Synthesis Of (E)-2-(3-phenyl-n-propylaminocarbonyl)-3-(3-isopropyl-4-hydroxy-5-(2-propylphenyl)acrylonitrile

A preferred method of synthesis of (E)-2-(3-phenyl-n-propylaminocarbonyl)-3-(3-isopropyl-4-hydroxy-5-(2-propylphenyl)acrylonitrile (Compound 5) is as follows: A mixture of 0.4g of 3,5-diisopropyl-6-hydroxy-benzaldehyde (for synthesis protocol, see Section 5.11(step 2)), 0.55g of N-3-phenyl-n-propyl cyanoacetamide (synthesized according to the protocol set forth in Gazit, et al., 1991, *J. Med. Chem.* 34(6):1896-1907) and 40 mg of  $\beta$ -alanine (in 20 ml ethanol) was refluxed for 5 hours. Workup (H<sub>2</sub>O, HCl, dichloromethane) gave an oil which crystallized on standing. Trituration with benzene-hexane gave 0.4 g light yellow solid having a melting point of 120°C. Another 0.55 g of solid was obtained from the mother liquid. The overall yield was 65%.

The product gave the following analytical data:  
NMR (CDCl<sub>3</sub>)  $\delta$  8.24 (1H, S, vinyl), 7.73 (2H, S), 7.23 (5H, m), 5.50 (1H, S, OH), 3.46 (2H, q, J=6.7 Hz, NH-CH<sub>2</sub>), 3.17 (2H, Septet, J=7.0 Hz), 2.71 (2H, t, J=6.7 Hz), 1.95 (2H, quintet, J=6.7 Hz), 1.30 (12H, d, J=7.0 Hz).

#### 5.6. Synthesis Of (E)-2-(benzylaminocarbonyl)-3-(3-iodo-4,5-dihydroxyphenyl)acrylonitrile

A preferred method of synthesis of (E)-2-(benzylaminocarbonyl)-3-(3-iodo-4,5-

dihydroxyphenyl)acrylonitrile (Compound 6) is as follows: A mixture of 0.4 g of 2-cyano-3-(4-hydroxy-3-iodo-5-methoxy)phenylacrylonitrile (which was prepared by condensation of 4-hydroxy-3-iodo-5-methoxybenzaldehyde with N-benzylcyanoacetamide) and 0.5ml of BBr<sub>3</sub> in 20ml dichloromethane was stirred for 2 hours at room temperature. Workup (H<sub>2</sub>O, ethyl acetate) gave 0.16 g of a yellow solid (41% yield) having a melting point of 220°C.

The product gave the following analytical data: NMR (acetone-d<sub>6</sub>)  $\delta$  8.05(1H, s, vinyl), 7.85(1H, d, J=2.1 Hz), 7.70(1H, d, J=2.1 Hz), 7.30(5H, m), 4.6(2H, s).

**5.7. Synthesis Of (E,E)-2-[[2-[[1-cyano-2(3,4-dihydroxyphenyl)ethenyl]carbonylamino]ethyl]aminocarbonyl-3-(3,4-dihydroxyphenyl)acrylonitrile**

A preferred method of synthesis of (E,E)-2-[[2-[[1-cyano-2(3,4-dihydroxyphenyl)ethenyl]carbonylamino]ethyl]aminocarbonyl-3-(3,4-dihydroxyphenyl)acrylonitrile (Compound 7) is as follows: A mixture of 0.7 g of 3,4-dihydroxybenzaldehyde, 0.5 g of N-(cyanomethylcarbonylamino-n-propyl)cyanoacetamide and 4 drops piperidine in 20 ml ethanol was refluxed for 4 hours. Water was added. Extraction with ethyl acetate, evaporation and trituration with ethanol-dichloromethane gave 0.34g (32% yield) of a yellow solid, having a melting point of 277°C.

**5.8. Synthesis Of (E,E)-2-[[4-[[1-cyano-2-(3,4-dihydroxyphenyl)ethenyl]carbonylamino]-n-butyl]aminocarbonyl-3-(3,4-dihydroxyphenyl)acrylonitrile**

A preferred method of synthesis of (E,E)-2-[[4-[[1-cyano-2-(3,4-dihydroxyphenyl)ethenyl]carbonylamino]-n-butyl]aminocarbonyl-3-(3,4-dihydroxyphenyl)acrylonitrile (Compound 8) is as

follows: The compound was prepared according to the protocol prescribed for the synthesis of compound above (Section 5.7, above). Following this protocol, a yellow solid (86% yield) resulted having a melting point of 283°C.

The product gave the following analytical data: NMR (acetone-d<sub>6</sub>)  $\delta$  8.25 (1H,5,vinyl), 7.23 (2H,5,H2,6).

5.9. Synthesis Of (E,E)-[2-[2-cyano-2-(3,4-dihydroxyphenylcarbonyl)ethenyl]sulfonyl]-3-(3,4-dihydroxyphenyl)acrylonitrile

A preferred method of synthesis of (E,E)-[2-[2-cyano-2-(3,4-dihydroxyphenylcarbonyl)ethenyl]sulfonyl]-3-(3,4-dihydroxyphenyl)acrylonitrile (Compound 9) is as follows: A mixture of 0.3 g of N-(2-cyanomethylcarbonylaminoethyl)cyanoacetamide and 0.55 g of 3,4-dihydroxy benzaldehyde with 30 mg  $\beta$ -alanine in 30 ml ethanol was refluxed for 6 hours and worked up. Evaporation of the ethyl acetate and trituration with acetone-benzene gave a 0.45 g (58% yield) of a yellow solid, having a melting point of 264°C (stains hands).

The product gave the following analytical data: MS m/e 312 (28%), 265 (24), 211 (100%), 185 (15%), 161 (44%), 160 (M-(compound structure), 80%), 157 (35), 129 (22), 114 (43) m/e. NMR (acetone-d<sub>6</sub>)  $\delta$  8.15 (2H,s,vinyl), 7.77 (2H,d,J=2.3 Hz, H2), 7.59 (2H,dd,J=8.4, 2.3 Hz, H6), 7.07 (2H,d,J=8.4 Hz, H5).

5.10. Synthesis Of (E)-3-(indol-5-yl)-2-(3,4-dihydroxyphenyl carbonyl)acrylonitrile

A preferred method of synthesis of (E)-3-(indol-5-yl)-2-(3,4-dihydroxyphenylcarbonyl)acrylonitrile (Compound 10) is as follows: A mixture of 130 mg of 5-formyl indole,

180 mg of cyanomethyl-(3,4-dihydroxyphenyl)ketone (prepared according to the protocol set forth at Gazit, et al., 1991, *J. Med. Chem.* 34:1896-1907) and 30 mg  $\beta$ -alanine was refluxed for 3 hours. Water was then added and the reaction extracted with ethyl acetate to give an oily solid containing some aldehyde. Chromatography gave 86 mg of a pure orange solid (28% yield) having a melting point of 185°C.

The product gave the following analytical data: MS m/e 304 (M+, 8%), 177 (29), 137 (C<sub>6</sub>H<sub>3</sub> (OH)<sub>2</sub> CO<sup>+</sup>, 100), 117 (12), 116 (indole+, 15), 109 (93). NMR (acetone-d<sub>6</sub>)  $\delta$  8.40 (1H, d, J=1.6 Hz, H<sub>4</sub>), 8.18 (1H, s, vinyl), 8.03 (1H, dd, J=8.6, 1.6 Hz, H<sub>6</sub>), 7.63 (1H, d, J=8.6 Hz, H<sub>5</sub>), 7.54-7.40 (3H, m, H<sub>3</sub>+H<sub>2</sub>, 6), 7.0 (1H, d J=8.6 Hz, HS), 6.69 (1H, d J=3.2 Hz, H<sub>2</sub>).

**5.11. Synthesis Of (E)-2-[3-phenyl-n-propylaminothiocarbonyl]-3-(3,5-diisopropyl-4-hydroxyphenyl)acrylonitrile**

A preferred method of synthesis of (E)-2-[3-phenyl-n-propylaminothiocarbonyl]-3-(3,5-diisopropyl-4-hydroxyphenyl)acrylonitrile (Compound 11) is as follows:

1. A mixture of 6.2 g of N-phenylpropylcyanoacetamide (synthesized according to the protocol set forth at Gazit, et al., 1991, *J. Med. Chem.* 34(6):1896-1907) and 15 g Lawesson reagent in 60 ml toluene was refluxed for 3 hours. Chromatography resulted in 1.5 g (22% yield) of N-phenylpropylcyanothioacetamide as a red solid. The product gave the following analytical data: NMR (CDCl<sub>3</sub>)  $\delta$  7.3 (5H, m), 3.81 (2H, s), 3.71 (2H, q, J=7.0 Hz), 2.74 (2H, t, J=7.0 Hz), 2.05 (2H, quintet, J=7.0 Hz).

2. A mixture of 18 g of 2,6-diisopropyl phenol and 1.8 g of hexamethylenetriamine (HMTA) in 60 ml



trifluoroacetic acid (TFA) was refluxed for 3.5 hours. Workup, chromatography and trituration with hexane gave 5.3 g (26% yield) of 3,8-diisopropyl-6-hydroxy-benzaldehyde white solid, having a melting point of 103°C. Analytical analysis of the product gave the following data: NMR (CDCl<sub>3</sub>)  $\delta$  9.87 (1H, S, CHO), 7.63 (2H, S), 3.19 (2H, septet, J=7.7 Hz), 1.30 (12H, d, J=7.7 Hz).

3. 0.6g of the compound of Step 1 (N-phenylpropyl cyanothioacetamide), 0.6g of the compound of Step 2 and 40 mg  $\beta$ -alanine in 40 ml ethanol were refluxed for 4 hours. Evaporation and chromatography gave 0.6g (50% yield) of Compound 11 as a viscous oil. NMR (CDCl<sub>3</sub>)  $\delta$  8.76 (1H, S, vinyl), 7.78 (2H, S, H<sub>2</sub>.6), 7.25 (5H, m), 5.60 (1H, S, OH), 3.90 (2H, q, J=7.0 Hz), 3.17 (2H, Septet, J=7.0 Hz), 2.76 (2H, t, J=7.0 Hz), 2.11 (2H, quintet, J=7.0 Hz), 1.29 (12H, d, J=7.0 Hz). MS m/e 407 (M+1, 55), 406 (M+, 70), 373 (M-CH<sub>3</sub>-H<sub>2</sub>O, 100), 363 (M-isopropyl, 72), 272 (M-NH(CH<sub>2</sub>)<sub>3</sub> Ph, 20), 259 (58), 230 (28), 91 (28).

5.12. Synthesis Of (E)-2-[1-cyano-2-(5-bromo-3,4-dihydroxyphenyl)ethenylsulfonyl]-3-(3-bromo-4,5-dihydroxy phenyl)acrylonitrile

A preferred method of synthesis of (E)-2-[1-cyano-2-(5-bromo-3,4-dihydroxyphenyl)ethenylsulfonyl]-3-(3-bromo-4,5-dihydroxy phenyl)acrylonitrile (Compound 12) is as follows: 10 ml of ethanol containing 230 mg of 5-bromo 3,4-dihydroxy benzaldehyde, 76 mg of diacetonitrile sulphone and 10 mg of  $\beta$ -alanine were refluxed for 5 hours. Cooling and filtering gave 220 mg (76% yield) of an orange solid having a melting point greater than 300°C.

The product gave the following analytical data:  
NMR (acetone- $d_6$ )  $\delta$  8.18(2H,S,vinyl), 7.90 (2H,d,J=1.6 Hz), 7.78 (2H,d,J=1.6 Hz).

5        5.13.        **Synthesis Of 2-[(4-iodophenyl)amino]-6,7-dimethyl quinoxaline**

A preferred method of synthesis of 2-[(4-iodophenyl)amino]-6,7-dimethyl quinoxaline (Compound 13) is as follows:

10        1.        A mixture of 2 g of 4,5-dimethyl-1,2-diaminobenzene and 1.5g of glyoxalic acid hydrate in 30 ml ethanol was refluxed for 2 hours. Cooling and filtering gave 1.2 g (46% yield) of a white solid, having a melting point of 263°C. The product gave the following analytical data:

15        NMR (DMSO  $d_6$ )  $\delta$  60:40 mixture.

major - 8.07 (1H,S), 7.55(1H,S), 7.06(1H,S), 2.30(6H,S).

minor - 8.02 (1H,S), 7.42(1H,S), 7.28(1H,S), 2.28(6H,S).

20        2.        A mixture of 1.1 g of the product of step 1 (6,7-dimethylquinoxalone), 1 ml phosphorous oxychloride ( $POCl_3$ ) and 1 ml dimethyl aniline in 20 ml toluene were refluxed for 2 hours. Workup ( $NH_3$ , dichloromethane) and chromatography gave 0.4 g (33% yield) of a white  
25        solid (2-chloro-6,7-dimethylquinoxaline), having a melting point 86°C. The product gave the following analytical data: NMR ( $CDCl_3$ )  $\delta$  8.68(1H,S, $H_2$ ), 7.85(1H,S), 7.76 (1H,S), 2.50(6H,S).

30        3.        A mixture of 210 mg of the product of step 2 (2-chloro-6,7-dimethylquinoxaline) and 0.8g of p-iodoaniline was heated in 10 ml ethanol at 100°C for 4 hours. Chromatography gave 245 mg (60% yield) of a light green solid having a melting point of 228°C.

35

The product gave the following analytical data:  
NMR(CDCl<sub>3</sub>)  $\delta$  8.32(1H,S), 7.67(1H,S), 7.64(1H,S), 7.68,  
7.56(4H,Abq, JAB=9.0 Hz).

5        5.14.        **Synthesis Of 2-(3,4-dihydroxyphenyl)-  
6,7-dimethyl-quinoxaline**

A method of synthesis of 2-(3,4-dihydroxyphenyl)-6,7-dimethyl-quinoxaline (Compound 14) is as follows: 1.4 g of 4,5-dimethyl 1,2-phenylenediamine and 1.9 g of  $\alpha$ -chloro 3,4-dihydroxy acetophenone in 15 ml dimethylsulfoxide were heated  
10        for 1.5 hours at 100°C. 80 ml of H<sub>2</sub>O was added and the suspension was left overnight at room temperature and filtered to give 2.5 g (67% yield) of a brown solid.

The product gave the following analytical data:  
15        NMR (acetone-d<sub>6</sub>)  $\delta$  9.28(1H,S,H<sub>2</sub>), 8.40(Br.S,OH),  
7.89(1H,d,J=2.2 Hz, H<sub>2</sub>'), 7.82 (2H,S,H<sub>5</sub>,8),  
7.72(1H,dd,J=8.3,2.2 Hz,H<sub>6</sub>'), 7.02 (1H,d,J=8.3  
Hz,H<sub>5</sub>'), 2.52 (6H,S,CH<sub>3</sub>). DMSO d<sub>6</sub> 9.30(1H,S,H<sub>2</sub>),  
7.81(2H,S,H<sub>5</sub>,8), 7.75 (1H,d,J=2.2 Hz,H<sub>2</sub>'),  
20        7.62(1H,dd,J=8.3,2.2 Hz,H<sub>6</sub>'), 6.90 (1H,d,J=8.3 Hz,  
H<sub>5</sub>'), 2.44 (6H,S,CH<sub>3</sub>).

A second method of synthesis of 2-(3,4-dihydroxy)phenyl-6,7-dimethyl-quinoxaline is as follows: 1 g and 1.9 g of the above reagents in 25 ml  
25        ethanol were refluxed 2 hours. Cooling and filtering gave 0.76 g (18% yield) of a deep yellow solid having a melting point of 278°C as the HCl salt.

5.15.        **Synthesis Of 4-(4-iodophenylamino)-6,7-dimethoxy quinazoline**

30        A preferred method of synthesis of the compound (Compound 15) is as follows:

1. A mixture of 7 g of 4,5-dimethoxy 2-aminobenzoic acid and 8 ml of formamide was heated for 2 hours at 170°C. Cold water was added and the solid  
35        filtered to give 0.9 g (12% yield) of a light-brown

solid (6,7-dimethoxyquinazolone), having a melting point of 308°C.

The product gave the following analytical data:  
NMR(DMSO-d<sub>6</sub>) δ 8.0(1H,S), 7.43(1H,S), 7.12(1H,S),  
3.89(3H,S), 3.85(3H,S).

2. 0.8 g of the compound of Step 1. 1 ml POCl<sub>3</sub> and 1 ml dimethylaniline in 20 ml toluene were refluxed for 3.5 hours. Workup and trituration with hexane gave 0.5 g of a light grey solid (57% yield), having a melting point of 188°C.

The product gave the following analytical data:  
NMR (CDCl<sub>3</sub>) δ 8.88(1H,S), 7.41(1H,S), 7.36(1H,S), 4.09(3H,S), 4.08(3H,S).

3. A mixture of 300 mg of 4-chloro-6,7-dimethoxyquinazoline and 300 mg of 3-iodoaniline in 10 ml ethanol was refluxed for 1 hour. Cooling and filtering gave 540 mg (93% yield) of a white solid as the HCl salt. The solid had a melting point of 278°C.

The product gave the following analytical data:  
NMR (DMSO-d<sub>6</sub>) δ 8.87(1H,S,H<sub>2</sub>), 8.27(1H,S), 8.13(1H,S), 7.8-7.66(2H,m), 7.33(2H,m), 4.02(3H,S), 4.0(3H,S).

#### 5.16. Synthesis Of 4-(3-hydroxyphenylamino)-6-methylquinazoline

A preferred method of synthesis of 4-(3-hydroxyphenylamino)-6-methylquinazoline (Compound 16) is as follows:

1. A mixture of .8 g of 5-methyl-2-aminobenzoic acid and 15 ml formamide was heated at 170°C for 1.5 hours. Water was added and the solid filtered to give 7.3 g (83% yield) of a brown-white solid (6-methylquinazolone) having a melting point of 268°C.

2. A mixture of 5 g of the compound of step 1, 5 ml of POCl<sub>3</sub> and 5 ml dimethyl aniline in 40 ml toluene were refluxed for 3.5 hours. Workup (NH<sub>3</sub>, H<sub>2</sub>O, ethyl acetate) gave a dark solid. Chromatography

yielded (29% yield) 1.61 g of a white solid (4-chloro-6-methyl-quinazoline) having a melting point of 98°C. The product gave the following analytical data: NMR (CDCl<sub>3</sub>) δ 9.0 (1H, s, H<sub>2</sub>), 8.04 (1H, d, J=2.0 Hz, H<sub>5</sub>), 7.96 (1H, d, J=8.8 Hz, H<sub>8</sub>), 7.80 (1H, dd, J=8.8, 2.0 Hz, H<sub>7</sub>), 2.62 (3H, s).

3. A mixture of 230 mg of the compound of step 2 and 145 mg of m-aminophenol in 10 ml ethanol was refluxed for 50 minutes. Cooling and filtering gave 300 mg (80% yield) of a light yellow solid, as the HCl salt. The product had a melting point of 262°C.

The product gave the following analytical data: (DMSO-d<sub>6</sub>) δ 8.89 (1H, s, H<sub>2</sub>), 8.72 (1H, s, H<sub>5</sub>), 7.90 (2H, ABq, J=8.0 Hz, H<sub>7,8</sub>), 7.2 (3H, m), 6.75 (1H, m), 2.55 (3H, s).

5.17. **Synthesis Of 2-(3,4-dichlorophenylamino)-6,7-dimethylquinoxaline**

A preferred method of synthesis of 2-(3,4-dichlorophenylamino)-6,7-dimethylquinoxaline (Compound 17) is as follows: A mixture of 150 mg of the compound of 2-chloro-6,7-dimethylquinoxaline (which may be synthesized according to the protocol at Section 5.13 (step 2)) and 0.7 g of 3,4-dichloroaniline was heated at 100°C for 3.5 hours. Chromatography gave 80 mg (33% yield) of a yellow brown solid, having a melting point of 229°C.

The product gave the following analytical data: NMR (CDCl<sub>3</sub>) δ 8.32 (1H, s), 8.16 (1H, d, J=2.4 Hz, H<sub>2'</sub>), 7.71 (1H, Br. s), 7.65 (1H, Br. s), 7.57 (1H, dd, J=2.4, 9.2 Hz, H<sub>6'</sub>), 7.43 (1H, d, J=9.2 Hz, H<sub>5'</sub>), 2.48 (3H, s), 2.46 (3H, s).

5.18. **Synthesis Of 4-(3-hydroxyphenylamino)-quinazoline**

A preferred method of synthesis of 4-(3-hydroxyphenylamino)-quinazoline (Compound 18) is as follows:

5 1. A mixture of 4.6 g of quinazolone, 5 ml of POCl<sub>3</sub>, and 5 ml of dimethylaniline in 50 ml toluene was refluxed for 3.5. hours. Workup (NH<sub>3</sub>, H<sub>2</sub>O and ethylacetate) yielded a green oil. Chromotography resulted in a light brown solid. Sublimation at 160°C (11 mm Hg) gave 1.48 g (29% yield) of a white solid  
10 (4-chloro-quinazoline) having a melting point of 72°C. This product gave the following analytical data: NMR (CDCl<sub>3</sub>) δ 9.05 (1H,S), 8.27 (1H,m), 8.1-7.9 (2H,m) 7.75 (1H,m).

15 2. A mixture of 0.37 g of the compound of step 1 and 0.24 g of m-hydroxyaniline in 10 ml ethanol was refluxed for 1 hour. Cooling and filtering gave 0.25 g (41% yield) of a light-yellow solid as the HCl salt. The solid turns light green on standing overnight.  
20 The product has melting point of 268°C.

The product gave the following analytical data: NMR (DMSO-d<sub>6</sub>) δ 8.99 (1H,S), 8.93(2H,S), 8.15-7.81 (3H,m), 7.31-7.12(3H,m), 6.77(1H,d,J=7.4 Hz).

5.19. **Synthesis Of 2-(n-propylamino)-3-chloroquinoxaline**

25 A preferred method of synthesis of 2-(n-propylamino)-3-chloroquinoxaline (Compound 19) is as follows: 0.8g of 2-phenyl-3,6,7-trimethyl quinoxaline 10ml dimethylsulfoxide was heated for 40 minutes at  
30 90°C. Workup and chromatography gave 40 mg (4% yield) of a white solid having a melting point of 148°C. The product gave the following analytical data: NMR (CDCl<sub>3</sub>) δ 8.03(1H,S), 7.90(1H,S), 7.60(5H,m), 6.96(1H,S,CHBr<sub>2</sub>), 2.52(6H,S).

35 Following the same protocol, 0.3 g (47% yield) of a white solid was obtained having a melting point of

150°C. The product gave the following analytical data: NMR(CDCl<sub>3</sub>) δ 11.30(1H,S,CHO), 8.05(1H,S), 7.96(1H,S), 7.67(2H,m), 7.55(3H,m), 2.55(6H,S).

5       **5.20.       Synthesis Of 4-[(4-methylphenyl)mercapto]quinazoline**

10       A preferred method of synthesis of the compound (Compound 20) is as follows: A mixture of 250 mg of 4-chloro-6-methyl-quinazoline, 180 mg of p-thiocresole and 100 mg of potassium hydroxide (KOH) are combined in 20 ml CH<sub>3</sub>CN and stirred 24 hours at room temperature. Workup (H<sub>2</sub>O, dichloromethane) and trituration with hexane gave 40 mg (10% yield) of a light blue solid having a melting point of 96°C.

15       The product gave the following analytical results: NMR (CDCl<sub>3</sub>) δ 8.81(1H,S), 7.96(1H,d,J=2.0 Hz,H5), 7.85(1H,d,J=9.0Hz, H8), 7.68(1H,dd,J=9.0, 2.0 Hz, H7), 7.51, 7.30(4H,ABq,JAB=8.2 Hz), 2.56(3H,S), 2.42(3H,S). MS m/e 266 (M<sup>+</sup>,40%), 265 (M-1,100%) m/e.

20       **5.21.       Synthesis Of 2-chloro-4-(3-chlorophenylamino)-6,7-dimethoxyquinazoline**

A preferred method of synthesis of 2-chloro-4-(3-chlorophenylamino)-6,7-dimethoxyquinazoline (Compound 21) is as follows:

25       1. A mixture of 8 g of 6,7-dimethoxy-2,4-quinazolinedione, 23 ml of POCl<sub>3</sub> and 10 ml of dimethylaniline in 30 ml toluene was refluxed for 5 hours. Workup (H<sub>2</sub>O, NH<sub>3</sub>, dichloromethane) and titration with hexane gave 7 g (75% yield) of a light green solid (2,4-dichloro-6,7-dimethoxy-quinazoline having a melting temperature of 156°C. The product gave the following analytical results: NMR (CDCl<sub>3</sub>) δ 7.36(1H,S), 7.28(1H,S), 4.07(3H,S), 4.06(3H,S).

30       2. A mixture of 2.6 g of the compound of step 1 and 1.3 g of m-chloroaniline in 20 ml ethanol was

refluxed for 1 hour. The mixture was then cooled and filtered to give 3.5 g (90% yield) pink-white solid.

Alternatively, a second method of synthesis using the free base was used. 3.4 g of the material form above was treated with  $\text{NH}_3\text{-H}_2\text{O}$  and extracted with ethyl acetate. Recrystallization from benzene-hexane gave 2.3 g (74% yield) of a white solid having a melting point of 222°C.

The product gave the following analytical data: NMR ( $\text{CDCl}_3$ )  $\delta$  7.74 (1H, t,  $J=2.2$  Hz,  $\text{H}_2'$ ), 7.63 (1H, m), 7.30 (1H, m), 7.16 (1H, s), 7.12 (1H, m), 6.98 (1H, t,  $J=8.0$  Hz), 4.0 (3H, s), 3.97 (3H, s).

5.22. **Synthesis Of (Z)-1-(2-chlorophenyl)-2-[2,2-dicyanoethenyl]hydrazine**

A preferred method of synthesis of (Z)-1-(2-chlorophenyl)-2-[2,2-dicyanoethenyl]hydrazine (Compound 22) is as follows: 2.4 g of sodium nitrite ( $\text{NaNO}_2$ ) was added to 4 g of m-chloro-aniline in 20 ml of diluted hydrochloric acid and 20 ml  $\text{H}_2\text{O}$  and then cooled in ice for approximately 0.5 hours. The mixture was then added into a solution of 2.2 g malononitrile, and 10 g potassium acetate in 100 ml ethanol. After 0.5 hours in the cold and 1 hour at room temperature the solid was filtered, washed with water and dried to give 2.4 g (37% yield) yellow solid.

The product gave the following analytical data: mp-170°C. NMR ( $\text{CDCl}_3$ )  $\delta$  7.4-7.2, m.



5.23.        **Synthesis Of 2-phenyl-1,4-diaza-anthracene**

A preferred method of synthesis of 2-phenyl-1,4-diaza-anthracene (Compound 23) is as follows: 20 ml of ethanol containing 0.47 grams of 2,3-diaminonaphthalene and 0.47 grams of phenyl gloxal hydrate were refluxed for 1.5 hour. Cooling and filtering gave 0.5g (65%) of a light brown solid with a melting point of 163°C.

The product gave the following analytical data: NMR (CDCl<sub>3</sub>): δ 9.38 (1H, l.c., H2), 8.71, 8.67 (2H, 2d, H5,10), 8.25, 8.10 (4H, AA'BB'm, H6-9), 7.58 (5H, m, Ph). MS m/e 256 (M+, 100%), 229 (M-CN, 12%), 126 (71%) m/e.

5.24.        **Synthesis Of N-(2,5-dihydroxylbenzyl)-4-methoxycarbonyl aniline**

A preferred method of synthesis of N-(2,5-dihydroxylbenzyl)-4-methoxycarbonylaniline (Compound 24) is as follows: 0.7 g of 2,5-dihydroxybenzaldehyde and 0.75 g of 3-aminomethylbenzoate in 40 ml of methanol were refluxed for 3 hours and cooled. 0.5 g of sodiumcyanoborohydride (NaCNBH<sub>4</sub>) was then added. After 12 hours at room temperature, workup (H<sub>2</sub>O, ethylacetate), and chromatography (silica gel, elution with 5% CH<sub>3</sub>OH in dichloromethane), 0.42 g (31% yield) of a light yellow solid was obtained.

The product gave the following analytical data: mp 175°C. NMR (acetone-d<sub>6</sub>) δ 7.78, 6.68 (4H, ABq, JAB=8.8 Hz), 6.74 (1H,d,J=3.0 Hz, H6), 6.72 (1H,d,J=8.5 Hz,H3), 6.55 (1H,d,J=8.5,3.0 Hz,H<sub>4</sub>), 4.34 (2H,s,CH<sub>2</sub>N), 3.76 (3H,s,COOCH<sub>3</sub>).

5.25.        **Synthesis Of N-(2-chlorophenyl)-2-cyano-2-(N-3-trifluorophenylaminocarbonyl)-thioacetamide**

N-(2-chlorophenyl)-2-cyano-2-(N-3-trifluorophenylamino-carbonyl)-thioacetamide (Compound 25) may be synthesized as follows:

5 680 mg of sodium ethoxide was added to a solution of 1.6 gram of N-3-trifluoromethylphenyl cyanoacetamide in 20 ml of tetrahydrofuran at 0°C. This mixture was stirred at 0°C for 1 hour and 1.7 g of 2-chlorophenylisothiocyanate in 5 ml of tetrahydrofuran was added dropwise. After addition, 10 the mixture was warmed to room temperature and heated at 50°C for 6 hours. Upon cooling, all the ethanol was removed and the resulting solid was resuspended in 10 ml of water. This was then added to 15 ml of 0.3 M sodium hydroxide solution, shaken vigorously and washed in 50 ml of ethyl ether. The aqueous layer was then acidified with 1 N hydrochloric acid to pH 1. The solid was then collected by suction filtration to produce 630 mg of N-2-chlorophenyl(2-cyano-2-N-3-trifluorophenylaminocarbonyl) thioacetamide.

20 5.26. Synthesis Of (E)-2-cyanomethylsufonyl-3-(3-bromo-4,5-dihydroxyphenyl)acrylonitrile

A preferred method of synthesis of (E)-2-cyanomethylsufonyl-3-(3-bromo-4,5-dihydroxyphenyl)acrylonitrile (Compound 26) is as 25 follows: A mixture of 500 mg of 5-bromo-3,4-dihydroxybenzaldehyde and 700 mg of sulfonyldiacetonitrile in 6 ml of ethanol was refluxed with a few drops of piperidine for 4 hours. Ethanol 30 was removed in a rotavap and the mixture worked up with ethyl acetate, diluted acid and brine. A portion of the crude was then purified by HPLC on a C-18 column to provide about 50 mg of (E)-2-cyanomethylsufonyl-3-(3-bromo-4,5-dihydroxyphenyl)acrylonitrile. 35

5.27.        Synthesis Of (1-benzyl-2-hydroxyphenyl)pyrrolido[3,4-b]3,4-dihydro-4-oxoquinazoline

A preferred method of synthesis of (1-benzyl-2-hydroxyphenyl)pyrrolido[3,4-b]3,4-dihydro-4-oxoquinazoline (Compound 27) is as follows: 0.01 mol (1.07 g) of benzylamine and 0.01 mol (1.22 g) of 4-hydroxybenzaldehyde were mixed together in 15 ml of ethanol and refluxed on a waterbath for 15 minutes. 0.01 mol (3.44 g) of 2-(1'-tosyloxyethyl)-quinazolin-4-one and one drop of pyridine were added and the mixture was refluxed for six hours. The resulting solution was evaporated and extracted with a 5% water solution of sodium bicarbonate. The remaining crystals were filtered off, washed with water and recrystallized from isopropanol. 3.40 g of compound was obtained (89% yield) having a melting point of 217-219°C.

The product ( $C_{24}H_{21}N_3O_2$ ) gave the following analytical data:

Elemental analysis [%] of Product

Calculated:            C: 75.18            H: 5.52            N: 10.96

Found:                C: 75.26            H: 5.47            N: 10.88

5.28.        Synthesis Of 2-(3-chlorophenylamino)-6,7-dimethylquinoxaline

A preferred method of synthesis of 2-(3-chlorophenylamino)-6,7-dimethylquinoxaline (Compound 28) is as follows: 200 mg of 2-chloro-6,7-dimethylquinoxaline and 700 mg m-chloro aniline were heated without solvent at 100°C for 2.5 hours. Chromatography gave 100 mg of a white solid (34% yield) having a melting point of 175°C.

The product gave the following data: NMR ( $CDCl_3$ ),  $\delta$  8.33 (1H, S), 7.97 (1H, m), 7.68 (1H, S), 7.62 (1H, S),

7.54 (1H,m), 7.27 (1H,m), 7.05 (1H,m), 2.45 (3H,S),  
2.43 (3H,S). MS m/e 285, 283 (M+, 29, 81%), 248 (m-Cl, 7).

5.29. **Synthesis Of (E)-2-(3,4-dihydroxybenzoyl)-3-dihydroxyphenyl)acrylonitrile.**

5 A preferred method of synthesis of (E)-2-(3,4-dihydroxybenzoyl)-3-dihydroxyphenyl)acrylonitrile (Compound 29) is as follows: Solid potassium cyanide (KCN) (3 g, 46 mmol) was added to 2-chloro-3',4'-  
10 dihydroxyacetophenone (6 g) in 30 ml of dimethylsulfoxide. The reaction mixture was stirred at 100°C for 2.5 hours. After cooling, 100 ml of 1 N hydrochloric acid was added, and the reaction mixture was extracted with ethyl acetate. Drying and  
15 evaporation gave a red oily solid which was purified by chromatography on silica gel. The first fractions eluted with 3% methanol in dichloromethane were evaporated and triturated with dichloromethane. Filtering gave 1 g (18% yield) of a light yellow solid  
20 having a melting point of 217°C.

The product gave the following analytical data:  
NMR (acetone-d6)  $\delta$  7.51 (2H,m,H2,6), 6.97 (1H,d,J=8.0 Hz, H5), 4.43 (2H,s,CH<sub>2</sub>CN).

5.30. **Synthesis Of 2-(4-bromophenylamino)-6,7-quinoxaline**

25 A preferred method of synthesis of 2-(4-bromophenylamino)-6,7-quinoxaline (Compound 30) is as follows: 200 mg of 2-chloro-6,7-dimethyl quinoxaline (synthesized according to the protocol at Section  
30 5.13, step 2) and 0.8g of p-bromoaniline were heated at 100° for 3.5 hours. Chromatography gave 25 mg (37% yield) of a light yellow solid, having a melting point of 235°C.

The product gave the following analytical data:  
35 NMR (CDCl<sub>3</sub>)  $\delta$  8.32 (1H,S), 7.68 (1H,S), 7.60 (1H,Br.S),

7.68, 7.48 (4H, ABq, JAB=8.8 Hz), 2.45 (3H, S),  
2.43 (3H, S).

5     **6.     EXAMPLE:   ELISA Assay To Measure Kinase Activity  
                    Of FLK-1 Receptor In FLK-1/NIH Cells**

      An ELISA assay was conducted to measure the  
kinase activity of the FLK-1 receptor and more  
specifically, the inhibition or activation of protein  
tyrosine kinase activity on the FLK-1 receptor.

10     **6.1. Materials And Methods**

**Materials.** The following reagents and  
supplies were used:

- a. Corning 96-well ELISA plates (Corning  
Catalog No. 25805-96);
- 15     b. Cappel Goat anti-rabbit IgG (catalog no.  
55641);
- c. PBS (Gibco Catalog No. 450-1300EB);
- d. TBSW Buffer (50 mM Tris (pH 7.2) m 150 mM  
NaCl and 0.1% Tween-20);
- 20     e. Ethanolamine stock (10% ethanolamine (pH  
7.0), stored at 4°C);
- f. HNTG buffer (20mM HEPES buffer (pH 7.5),  
150mM NaCl, 0.2% Triton X-100, and 10% Glycerol);
- g. EDTA (0.5 M (pH 7.0) as a 100X stock);
- 25     h. Sodium Ortho Vanadate (0.5 M as a 100X  
stock);
- i. Sodium pyro phosphate (0.2M as a 100X  
stock);
- j. NUNC 96 well V bottom polypropylene plates  
30     (Appplied Scientific Catalog No. AS-72092);
- k. NIH3T3C7#3 Cells (FLK-1 infected cells);
- l. DMEM with 1X high glucose L Glutamine  
(catalog No. 11965-050);
- m. FBS, Gibco (catalog no. 16000-028);
- 35     n. L-glutamine, Gibco (catalog no. 25030-016);

o. VEGF, PeproTech, Inc. (catalog no. 100-20) (kept as 1 ug/100 ul stock in Milli-Q dH<sub>2</sub>O and stored at -20°C;

5 p. Affinity purified anti-flk-1 antiserum, Enzymology Lab, Sugen, Inc.;

q. UB40 monoclonal antibody specific for phosphotyrosine, Enzymology Lab, Sugen, Inc.;

10 r. EIA grade Goat anti-mouse IgG-POD (BioRad catalog no. 172-1011);

s. 2,2-azino-bis(3-ethylbenz-thiazoline-6-sulfonic acid (ABTS) solution (100mM citric acid (anhydrous), 250 mM Na<sub>2</sub>HPO<sub>4</sub> (pH 4.0), 0.5 mg/ml ABTS (Sigma catalog no. A-1888)), solution should be stored in dark at 4°C until ready for use;

15 t. H<sub>2</sub>O<sub>2</sub> (30% solution) (Fisher catalog no. H325);

u. ABTS/H<sub>2</sub>O<sub>2</sub> (15ml ABTS solution, 2 ul H<sub>2</sub>O<sub>2</sub>) prepared 5 minutes before use and left at room temperature;

20 v. 0.2 M HCl stock in H<sub>2</sub>O;

w. dimethylsulfoxide (100%) (Sigma Catalog No. D-8418); and

y. Trypsin-EDTA (Gibco BRL Catalog No. 25200-049).

25 **Protocol.** The following protocol was used to conduct the ELISA Assay:

1. Coat Corning 96-well elisa plates with 1.0ug per well Cappel Anti-rabbit IgG antibody in 0.1M Na<sub>2</sub>CO<sub>3</sub> pH 9.6. Bring final volume to 150 ul per well. Coat plates overnight at 4 C. Plates can be kept up to two weeks when stored at 4°C.

30 2. Grow cells in 30 ml of Growth media (9DMEM, 2.0mM L-Glutamine, 10% FBS) until confluent in 150cm tissue culture dishes at 37°C, 5% CO<sub>2</sub>.

35

3. Harvest cells by trypsinization and seed in Corning 25850 polystyrene 96-well roundbottom cell plates, 25,000 cells/well in 200uL of growth media.
- 5 4. Grow cells at least one day at 37°C, 5% CO<sub>2</sub>.
5. Wash cells with D-PBS 1X.
6. Add 200ul/well of starvation media. (DMEM, 2.0mM l-Glutamine, 0.1% FBS). Incubate overnight at 37°C, 5% CO<sub>2</sub>.
- 10 7. Dilute Compounds/Extracts 1:20 in polypropylene 96 well plates using starvation media. Dilute dimethylsulfoxide 1:20 for use in control wells.
- 15 8. Remove starvation media from 96 well cell culture plates and add 162 ul of fresh starvation media to each well.
9. Add 18ul of 1:20 diluted Compound/Extract dilution (from step #7) to each well plus the 1:20 dimethylsulfoxide dilution to the control wells (+/- VEGF), for a final dilution of 1:200 after cell stimulation. Final dimethylsulfoxide is 0.5 %. Incubate the plate at 37°C, 5% CO<sub>2</sub> for two hours.
- 20 10. Remove unbound antibody from Elisa plates by inverting plate to remove liquid. Wash 3 times with TBSW + 0.5% Ethanolamine, pH 7.0. Pat the plate on a paper towel to remove excess liquid and bubbles.
- 25 11. Block plates with TBSW + 0.5% Ethanolamine, pH 7.0. 150 ul per well. Incubate plate thirty minutes while shaking on a microtiter plate shaker.
- 30 12. Wash plate 3 times as described in step 10.
- 35

13. Add 0.5ug/well affinity purified anti-flk-1 polyclonal rabbit antiserum. Bring final volume to 150ul/well with TBSW + 0.5% Ethanolamine pH 7.0. Incubate plate for thirty minutes while shaking.
14. Add 180  $\mu$ l starvation medium to the cells and stimulate cells with 20ul/well 10.0mM Sodium Ortho Vanadate and 500 ng/ml VEGF (resulting in a final concentration of 1.0mM Sodium Ortho Vanadate and 50ng/ml VEGF per well) for eight minutes at 37°C, 5% CO<sub>2</sub>. Negative control wells receive only starvation medium.
15. After eight minutes, media are removed from the cells and washed one time with 200ul/well PBS.
16. Lyse cells in 150ul/well HNTG while shaking at room temperature for five minutes. HNTG formulation includes sodium ortho vanadate, sodium pyro phosphate and EDTA.
17. Wash Elisa plate three times as described in step 10.
18. Transfer cell lysates from the cell plate to elisa plate and incubate while shaking for two hours. To transfer cell lysate pipette up and down while scrapping the wells.
19. Wash plate three times as described in step 10.
20. Incubate Elisa plate with 0.02ug/well UB40 in TBSW + 05% ethanolamine. Bring final volume to 150ul/well. Incubate while shaking for 30 minutes.
21. Wash plate three times as described in step 10.
22. Incubate elisa plate with 1:10,000 diluted EIA grade Goat anti-mouse IgG conjugated horseradish peroxidase in TBSW + 0.5% ethanolamine, pH



7.0. Bring final volume to 150ul/well. Incubate while shaking for thirty minutes.

23. Wash plate as described in step 10.

24. Add 100 ul of ABTS/H2O2 solution to well. Incubate ten minutes while shaking.

25. Add 100 ul of 0.2 M HCL for 0.1 M HCL final to stop the colordevelopment reaction. Shake 1 minute at room temperature. Remove bubbles with slow stream of air and read the ELISA plate in an ELISA plate reader at 410 nm.

### 6.2. Experimental Results

The results obtained for the tested compounds of the present invention are set forth at Table 2.

TABLE 2  
ELISA In Vitro Assay Results

FLK-1R ELISA ASSAY RESULTS			
Compound	IC50 (uM)	Compound	IC50 (uM)
7	37.5	31	3.3
8	10.8	31	33.9
1	0.8	32	18.0
9	27.4	12	7.1
3	4.9	4	8.9
5	0.7	6	21.2
11	9.8	33	48.4
26	11	34	1.8
35	0.7	23	4.4
14	9.3	19	25.6
36	34.6	28	16.0
37	20.3	17	15.0
13	10.4	18	34.60
16	35.1	20	17.8

FLK-1R ELISA ASSAY RESULTS			
Compound	IC50 (uM)	Compound	IC50 (uM)
38	29.2	21	9.9
27	3.4	25	8.5
39	10	10	17.7
24	28.9	40	54.0
22	2.3	41	>50
42	>50	43	>50
44	>50	45	>50
46	>50	47	~50
48	>50	49	>50
50	>50	51	>50
52	>50	53	>50
54	~50	55	>50
56	>50	57	12.3
58	>50	59	>50
60	0.3	61	>50
22	2.3	62	>50
63	17.0	64	3.7
65	14.3		

## 7. EXAMPLES: Effect Of Compounds In In Vivo Studies

The ability of one of the compounds of the present invention, leflunomide (Compound 40), to inhibit ovarian, melanoma, prostate, lung and mammary tumor cell lines established as SC xenografts was examined. These studies were conducted using doses ranging from 12 to 20 mg/kg/day.

### 7.1: Materials And Methods.

The tumor cells were implanted subcutaneously into the indicated strains of mice.

Treatment was initiated on day 1 post implantation unless otherwise indicated (e.g. treatment of the SCID mouse related to the A375 melanoma cell line began on Day 9). Eight (8) to ten (10) mice comprised each test group.

Specifically:

Animals. Female athymic mice (BALB/c, nu/nu), BALB/c mice, Wistar rats and Fisher 344 rats were obtained from Simonsen Laboratories (Gilroy, CA). Female A/I mice were obtained from Jackson Laboratory (Bar Harbor, ME). DA rats were obtained from B&K Universal, Inc. (Fremont, CA). Athymic R/Nu rats, DBA/2N mice, and BALB/c mice were obtained from Harlan Sprague Dawley (Indianapolis, IN). Female C57BL/6 mice were obtained from Taconic (Germantown, NY). All animals were maintained under clean-room conditions in Micro-isolator cages with Alpha-dri bedding. They received sterile rodent chow and water *ad libitum*.

All procedures were conducted in accordance with the NIH Guide for the Care and Use Of Laboratory Animals.

*Subcutaneous Xenograft Model.* Cell lines were grown in appropriate medium as described (See Section 6). Cells were harvested at or near confluency with 0.05% Trypsin-EDTA and pelleted at 450 x g for 10 min. Pellets were resuspended in sterile PBS or media (without FBS) to a suitable concentration indicated in the Figure legends and the cells were implanted into the hindflank of mice. Tumor growth was measured over 3 to 6 weeks using venier calipers tumor volumes were calculated as a product of length x width x height unless otherwise indicated. P values were calculated using the Students' t-test. sul01 in 50 - 100 uL excipient (dimethylsulfoxide, PBTE, PBTE6C:D5W, or

PBTE:D5W) was delivered by IP injection at concentrations indicated in the Figure legends.

5        *Intracerebral Xenograft Model.* For the mouse IC model, rat C6 glioma cells were harvested and  
suspended in sterile PBS at a concentration of  $2.5 \times 10^7$  cells/ml and placed on ice. Cells were implanted  
10 into BALB/c, nu/nu mice in the following manner: the frontoparietal scalps of mice were shaved with animal  
clippers if necessary before swabbing with 70%  
ethanol. Animals were anesthetized with isofluorane  
and the needle was inserted through the skull into the  
left hemisphere of the brain. Cells were dispensed  
15 from Hamilton Gas-tight Syringes using 30 ga 1/2 inch  
needles fitted with sleeves that allowed only a 3 mm  
penetration. A repeater dispenser was used for  
accurate delivery of 4 uL of cell suspension. Animals  
were monitored daily for well-being and were  
sacrificed when they had a weight loss of about 40%  
and/or showed neurological symptoms.

20        For the rat IC model, rats (Wistar, Sprague Dawley, Fisher 344, or athymic R/Nu; approximately 200 g) were anesthetized by an IP injection of 100 mg/kg  
Ketaset (ketamine hydrochloride; Aveco, Fort Dodge,  
25 Iowa) and 5 mg/kg Rompun (xylazine, 2% solution; Bayer, Germany). After onset of anesthesia, the scalp  
was shaved and the animal was oriented in a  
stereotaxic apparatus (Stoelting, Wood Dale, IL). The  
skin at the incision site was cleaned 3 times with  
alternating swabs of 70% ethanol and 10% Povidone-  
30 Iodine. A median 1.0 - 1.5 cm incision was made in  
the scalp using a sterile surgical blade. The skin  
was detached slightly and pulled to the sides to  
expose the sutures on the skull surface. A dental  
drill (Stopiting, Wood Dale, IL) was used to make a  
35 small (1-2 mm diameter) burrhole in the skull

approximately 1 mm anterior and 2 mm lateral to the bregma. The cell suspension was drawn into a 50 uL Hamilton syringe fitted with a 23 or 25g a standard bevel needle. The syringe was oriented in the burrhole at the level of the arachnoidea and lowered until the tip of the needle was 3 mm deep into the brain structure, where the cell suspension was slowly injected. After cells were injected, the needle was left in the burrhole for 1-2 minutes to allow for complete delivery of the cells. The skull was cleaned and the skin was closed with 2 to 3 sutures. Animals were observed for recovery from surgery and anesthesia. Throughout the experiment, animals were observed at least twice each day for development of symptoms associated with progression of intracerebral tumor. Animals displaying advanced symptoms (leaning, loss of balance, dehydration, loss of appetite, loss of coordination, cessation of grooming activities, and/or significant weight loss) were humanely sacrificed and the organs and tissues of interest were resected.

*Intraperitoneal Model.* Cell lines were grown in the appropriate media. Cells were harvested and washed in sterile PBS or medium without FBS, resuspended to a suitable concentration, and injected into the IP cavity of mice of the appropriate strain. Mice were observed daily for the occurrence of ascites formation. Individual animals were sacrificed when they presented with a weight gain of 40%, or when the IP tumor burden began to cause undue stress and pain to the animal.

*Immunohistochemistry.* Acetone-fixed, 5 um frozen tissue sections untreated xenograft tumors derived from human, rat, or murine tumor cells were analyzed by immunohistochemistry using highly specific receptor

antibodies. Briefly, non-specific binding sites were blocked with 10% normal goat serum prior to the application of the primary antibody. Appropriate antibody concentrations were used to achieve the desired sensitivity and specificity (rabbit anti-human PDGF-B receptor 1:400, and affinity purified rabbit anti-mouse FLK-1 5.5 ug/ml). Tissue sections known to contain the protein of interest served as positive controls. Appropriate negative controls of normal rabbit IgG and mouse anti-chicken IgG of the same protein concentration and isotype as the primary antibodies were used. The detection method was a three-step indirect procedure and consisted of the primary antibody bound to a biotin labeled secondary antibody (goat anti-rabbit IgG 1:500) followed by streptavidin conjugated horseradish peroxidase. Diaminobenzidine/0.03%hydrogen peroxide (1:200) (0.05%) was used as the chromogen/substrate. Tissue sections were counterstained with hematoxylin, dehydrated through ascending grades of ethanol, cleared in Xylene Substitute, and coverslipped with Permunt for microscopic evaluation. A "+" to "+++" grading system was used to identify the overall intensity of the expression. One plus ("+") reflects low intensity. Two pluses ("++") relates to medium intensity and three pluses ("+++") relates to high intensity. "T" is used in Table 4, below, to designate a tumor cell-specific staining reaction. "V" is used in the Table 4, below, to indicate a vascular endothelial cell-specific staining reaction. "\*" is used above to indicate that the analysis was carried out using cytopspins from the indicated cell lines. "NS" refers to "not significant" and "NT" refers to "not tested."

### 7.2. Experimental Results.

Acetone-fixed frozen sections from xenograft tumors derived from human or murine tumor cell lines were analyzed by immunohistochemistry (IHC), as discussed above, using highly specific receptor antibodies to determine which tumor cell lines expressed the FLK-1 receptor. Specifically, the antibodies were obtained according to the procedure set forth at United States Application Serial No. 08/193,829. The results of the IHC analysis is set forth at Table 4:

TABLE 4  
IHC Analysis Of Tumors

Tumor	FLK-1	% Inhibition
C6	++ (V)	>95%
SKOV3T	NT	>95%
D1B	NT	95%
SF763T	NT	85%
U87MG	NT	75%
L1210	NT	75%
PC-3	+ / ++ (V)	71%
SF767T	NT	70%
U118T	NT	57%
Calu-6	++ / +++ (V)	64%
U373MG	NT	54%
PA-1	++ (V)	53%
A375	++ / +++ (V)	53%
A431	-	NS
MCF7	-	NS
A549	NT	NS

Tumor	FLK-1	% Inhibition
MCF7/HER2	-	NT
SKOV3	NT	NT

The *in vivo* experiments described above were then performed using the tumor cell lines which expressed FLK-1. The results were obtained in the above-described *in vivo* experiments are set forth at Table 4.

TABLE 5  
Effect Of Compound 40 On Tumor Growth In Vivo

Tumor Type	Cell Line	Strain	Dose mg/kg/day	% Inhibition (day)	P <
ovarian	PA-1	nu/nu	20	53 (36)	0.04
melanoma	A375	nu/nu	20	53 (31)	0.03
melanoma	A375	SCID (day 9)	15	53 (31)	0.002
prostate	PC-3	nu/nu	20	71 (45)	0.01
prostate	PC-3	SCID (day 15)	12	47 (36)	0.001
lung	Calu-6	nu/nu	20	64 (28)	0.0001

These studies show a significant inhibition of tumor growth in immunocompetent animals treated with Compound 40. Specifically, as set forth above, Compound 40 effectively inhibited the growth of human ovarian (PA-1), human melanoma (A375), human prostate (PC-3), and human lung (Calu-6).

The studies were repeated to test additional compounds, including Compound 23. Compound 23 exhibited 41% inhibition at 20 mg/kg/day. More



specifically, the results observed are set forth at Table 5.

TABLE 6  
Effect Of Compounds In Vivo

Compound	CALU-6 mg/kg/day
4	no significant inhibition @ 20
23	41% inhibition @ 20
12	no detectable inhibition @ 20
2	no detectable inhibition at 20 or 40
40	63.9% inhibition @ 20

The apparent lack of inhibition for some of the tested compounds does not necessarily indicate lack of activity (inhibition of angiogenesis and/or vasculogenesis) and may be explained by, for example, the half-life of the compound tested in vivo or the dose administered in the experiments.


The present invention is not to be limited in scope by the exemplified embodiments which are intended as illustrations of single aspects of the invention, and any clones, DNA or amino acid sequences which are functionally equivalent are within the scope of the invention. Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description and accompanying drawings. Such modifications are intended to fall within the scope of the appended claims.

All references cited herein are hereby incorporated by reference in their entirety.

CLAIMS

What Is Claimed:


1. A compound having the formula:



and pharmaceutically acceptable salts thereof,  
wherein:

15  $R_1$  is isopropyl, t-butyl, Br, OH or methyl,  
 $R_2$  is OH,  
 $R_3$  is 2-propyl, t-butyl, OH, H or methyl, and  
 $R_4$  is (E) 1-cyano-2-[(3,5-diisopropyl-4-  
hydroxy)phenyl]ethenyl-sulfonyl, aminothiocarbonyl,  
20 cyanomethylsulfonyl, (3-amino-4-cyano)pyrazo-5-yl, (1-  
phenyl)-n-propylaminothiocarbonyl, (E) [[[4-[1-cyano-  
2(3,4-dihydroxy)phenyl]ethenyl]carbonylamino]-n-  
butyl]aminocarbonyl, 2[[2-cyano-1-(3,4-  
dihydroxy)phenyl]ethylenyl]]sulfonyl or  
25 hydroxycarbonyl.

2. A compound having the formula:



and pharmaceutically acceptable salts thereof,  
wherein:


35  $R_1$  is  $\text{CH}_3$  or H,

$R_2$  is H,

$R_3$  is H or chloro, and

$R_4$  is (4-methylphenyl)mecapto or (3-hydroxyphenyl)amino.

3. A compound having the formula:



D.


and pharmaceutically acceptable salts thereof, wherein:

$R_1$  is 4-hydroxy-phenyl.

$R_2$  is benzyl, and

$R_3$  is  $CH_3$ .

4. A compound having the formula:



E.


and pharmaceutically acceptable salts thereof, wherein:

$R_1$  is (3-trifluormethyl)phenyl and

$R_2$  is (2-chlorophenyl)aminothiocarbonyl.

5. A compound having the formula:

5 F.




10

and pharmaceutically acceptable salts thereof,  
wherein:

- 15  $R_1$  is O,  
 $R_2$  is (3,4-dihydroxy)phenyl, and  
 $R_3$ ,  $R_4$  and  $R_5$  are H.

6. A compound having the formula:

20 G.



20


and pharmaceutically acceptable salts thereof,  
wherein:

- 25  $R_1$  is 4-(1-nitro)thiophene or indol-3-yl and  
 $R_2$  is aminothiocarbonyl or (3-amino-4-cyano)pyrazol-5-yl.

7. A compound having the formula:

30

J.




35

and pharmaceutically acceptable salts thereof, wherein  $R_1$  is (3-chloro)phenylamino.

8. A pharmaceutical composition comprising a compound having the formula:

A.



and pharmaceutically acceptable salts thereof, wherein:

$R_1$  is isopropyl, t-butyl, I, Br, OH or methyl,

$R_2$  is OH,

$R_3$  is 2-propyl, t-butyl, OH, H or methyl, and

$R_4$  is (1-phenyl)-n-propylaminocarbonyl, (E)

1-cyano-2-[(3,5-diisopropyl-4-hydroxy)phenyl]ethenyl-sulfonyl,

aminothiocarbonyl, cyanomethylsulfonyl, (3-amino-4-cyano)pyrazo-5-yl, phenyl-n-

propylaminocarbonyl, (E) 1-cyano-2-[(5-bromo-3,4-dihydroxy)phenyl]ethenylsulfonyl, (1-phenyl)-n-

propylaminothiocarbonyl, cyano, (E) [[[4-[1-cyano-2(3,4-


dihydroxy)phenyl]ethenyl]carbonylamino]-n-butyl]aminocarbonyl, benzylaminocarbonyl, 2[[2-

cyano-1-(3,4-dihydroxy)phenyl]ethylenyl]]sulfonyl, [(3,4-dihydroxy)phenyl]carbonyl, (E) [[[4-[1-cyano-

2(3,4-dihydroxy)phenyl]ethenyl]carbonylamino]-ethyl]aminocarbonyl or hydroxycarbonyl;

-OR-

B.



and pharmaceutically acceptable salts thereof,  
wherein:

$R_1$  is  $\text{CH}_3$  or H,

$R_2$  is  $\text{CH}_3$  or H, or alternatively,


$R_1$  and  $R_2$  form a phenyl ring ( $\text{CHCHCHCH}$ ),

$R_3$  is H or formyl or chloro, and

$R_4$  is phenyl, (3,4-dihydroxy)phenyl, (4-iodophenyl)amino, (3,4-dichlorophenyl)amino, (3-chlorophenyl)amino, (4-bromophenyl)amino or n-propylamino;

-OR-

C.



and pharmaceutically acceptable salts thereof,  
wherein:


$R_1$  is  $\text{OCH}_3$ ,  $\text{CH}_3$  or H,

$R_2$  is  $\text{OCH}_3$  or H,

$R_3$  is H or chloro, and

$R_4$  is (3-chlorophenyl)amino, (4-methylphenyl)mecapto, (4-iodophenyl)amino or (3-hydroxyphenyl)amino;

-OR-




and pharmaceutically acceptable salts thereof,  
wherein:

$R_1$  is 4-hydroxy-phenyl

$R_2$  is benzyl, and

$R_3$  is  $\text{CH}_3$ ;

-OR-




and pharmaceutically acceptable salts thereof,  
wherein:

$R_1$  is (3-trifluormethyl)phenyl and

$R_2$  is (2-chlorophenyl)aminothiocarbonyl;

-OR-

5 F.



10

and pharmaceutically acceptable salts thereof,  
wherein:

$R_1$  is O,

$R_2$  is (3,4-dihydroxy)phenyl, and


15

$R_3$ ,  $R_4$  and  $R_5$  are H;

-OR-

G.

20



and pharmaceutically acceptable salts thereof,  
wherein:

25

$R_1$  is 4-(1-nitro)thiophene or indol-3-yl or  
indol-5-yl and

$R_2$  is aminothiocarbonyl, (3-amino-4-  
cyano)pyrazol-5-yl or (3,4-  
dihydroxyphenyl)carbonyl;

30


-OR-

35



H.

5



10

and pharmaceutically acceptable salts thereof,  
wherein:

R<sub>1</sub> is 2,5-dihydroxybenzyl,

R<sub>2</sub> is H,


R<sub>3</sub> is methoxycarbonyl, and

15

R<sub>4</sub> is H;

I.

20



25

and pharmaceutically acceptable salts thereof,  
wherein:

R<sub>1</sub> is H,

R<sub>2</sub> is (4-trifluoromethyl)phenyl, and


R<sub>3</sub> is methyl;

30

-OR-

35

J.



5


and pharmaceutically acceptable salts thereof,  
 wherein  $R_1$  is (3-chloro)phenylamino;  
 and a pharmaceutical excipient.

10

9. A method for modulating vasculogenesis  
 and/or angiogenesis comprising the step of  
 administering an effective amount of a compound having  
 the formula:

15

A.



20

and pharmaceutically acceptable salts thereof,  
 wherein:

25

$R_1$  is isopropyl, t-butyl, I, Br, OH or  
 methyl,

$R_2$  is OH,

$R_3$  is 2-propyl, t-butyl, OH, H or methyl, and

$R_4$  is (1-phenyl)-n-propylaminocarbonyl, (E)

30

1-cyano-2-[(3,5-diisopropyl-4-

hydroxy)phenyl]ethenyl-sulfonyl,

aminothiocabonyl, cyanomethylsulfonyl, (3-amino-

4-cyano)pyrazo-5-yl, phenyl-n-

propylaminocarbonyl, (E) 1-cyano-2-[(5-bromo-3,4-

35


dihydroxy)phenyl]ethenylsulfonyl, (1-phenyl)-n-

propylaminothiocabonyl, cyano, (E) [[[4-[1-

cyano-2 (3,4-  
 dihydroxy)phenyl]ethenyl]carbonylamino]-n-  
 butyl]aminocarbonyl, benzylaminocarbonyl, 2[[2-  
 cyano-1- (3,4-  
 dihydroxy)phenyl]ethylenyl]]sulfonyl, [(3,4-  
 dihydroxy)phenyl]carbonyl, (E) [[[4- [1-cyano-  
 2 (3,4-dihydroxy)phenyl]ethenyl]carbonylamino]-  
 ethyl]aminocarbonyl or hydroxycarbonyl;

-OR-

B.




and pharmaceutically acceptable salts thereof,  
 wherein:

R<sub>1</sub> is CH<sub>3</sub> or H,  
 R<sub>2</sub> is CH<sub>3</sub> or H, or alternatively,  
 R<sub>1</sub> and R<sub>2</sub> form a phenyl ring (CHCHCHCH),  
 R<sub>3</sub> is H or formyl or chloro, and  
 R<sub>4</sub> is phenyl, (3,4-dihydroxy)phenyl, (4-  
 iodophenyl)amino, (3,4-dichlorophenyl)amino, (3-  
 chlorophenyl)amino, (4-bromophenyl)amino or n-  
 propylamino;

-OR-

C.



and pharmaceutically acceptable salts thereof,  
wherein:

$R_1$  is  $\text{OCH}_3$ ,  $\text{CH}_3$  or  $\text{H}$ ,


$R_2$  is  $\text{OCH}_3$  or  $\text{H}$ ,

$R_3$  is  $\text{H}$  or chloro, and

$R_4$  is (3-chlorophenyl)amino, (4-methylphenyl)mecapto, (4-iodophenyl)amino or (3-hydroxyphenyl)amino;

-OR-

D.



and pharmaceutically acceptable salts thereof,  
wherein:

$R_1$  is 4-hydroxy-phenyl


$R_2$  is benzyl, and

$R_3$  is  $\text{CH}_3$ ;

-OR-

5

E.



10

and pharmaceutically acceptable salts thereof,  
wherein:

$R_1$  is (3-trifluormethyl)phenyl and


$R_2$  is (2-chlorophenyl)aminothiocarbonyl;

-OR-

15

F.

20



25

and pharmaceutically acceptable salts thereof,  
wherein:

$R_1$  is O,

$R_2$  is (3,4-dihydroxy)phenyl, and


$R_3$ ,  $R_4$  and  $R_5$  are H;

-OR-

30

35

G.




and pharmaceutically acceptable salts thereof,  
wherein:

$R_1$  is 4-(1-nitro)thiophene or indol-3-yl or  
indol-5-yl and

$R_2$  is aminothiocarbonyl, (3-amino-4-  
cyano)pyrazol-5-yl or (3,4-  
dihydroxyphenyl)carbonyl;

-OR-

H.



and pharmaceutically acceptable salts thereof,  
wherein:

$R_1$  is 2,5-dihydroxylbenzyl,


$R_2$  is H,

$R_3$  is methoxycarbonyl, and

$R_4$  is H;

-OR-

I.



and pharmaceutically acceptable salts thereof,  
wherein:


$R_1$  is H,

$R_2$  is (4-trifluoromethyl)phenyl, and

$R_3$  is methyl;

-OR-

J.



and pharmaceutically acceptable salts thereof,  
wherein  $R_1$  is (3-chloro)phenylamino.

10. A method for treating diseases related to  
vasculogenesis and/or angiogenesis comprising  
administration of an effective amount of the  
composition of Claim 8.

11. A method of Claim 10 wherein said disease is  
selected from the group comprising diabetes, diabetic  
retinopathy, hemangioma, glioma, melanoma, Kaposi's  
sarcoma and ovarian, breast, lung, pancreatic,  
prostate, colon and epidermoid cancer.

12. A method for inhibiting solid cell tumor  
growth comprising administration of an effective amount  
of the composition of Claim 8.

13. A method for screening compounds for the ability to regulate angiogenesis and/or vasculogenesis comprising:

- 5 (a) stimulating target cells by concurrent introduction of sodium ortho vandate and a FLK-1 specific ligand;
- 10 (b) contacting (i) a lysate of the stimulated target cell which was exposed to the test compound with (ii) an anchoring molecule specific for a protein substrate which is phosphorylated as a result of signal transduction in the target cell, under conditions and for a time sufficient to allow binding of the protein substrate to the anchoring molecule; and
- 15 (c) detecting phosphotyrosine residues on any protein substrate bound to the anchoring molecule, in which differences in the detection of phosphotyrosine residues on the immobilized protein substrate derived from lysates of the target cells which were exposed to the test substance as compared to that of immobilized protein substrate derived from control target cells which were not exposed to the test substance, indicate that the test substance modulates the activity of the tyrosine kinase directly or indirectly involved in a signal transduction.
- 20
- 25
- 30

14. An assay for screening compounds for the ability to regulate angiogenesis and/or vasculogenesis comprising:

35



- 5
- 10
- 15
- 20
- 25
- 30
- 35
- (a) stimulating target cells by concurrent introduction of sodium ortho vandate and a FLK-1 specific ligand;
  - (b) contacting (i) a lysate of the stimulated target cell which was exposed to the test compound with (ii) an anchoring molecule specific for a protein substrate which is phosphorylated as a result of signal transduction in the target cell, under conditions and for a time sufficient to allow binding of the protein substrate to the anchoring molecule; and
  - (c) detecting phosphotyrosine residues on any protein substrate bound to the anchoring molecule, in which differences in the detection of phosphotyrosine residues on the immobilized protein substrate derived from lysates of the target cells which were exposed to the test substance as compared to that of immobilized protein substrate derived from control target cells which were not exposed to the test substance, indicate that the test substance modulates the activity of the tyrosine kinase directly or indirectly involved in a signal transduction.

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US95/01751

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/7.21; 436/503; 514/94, 249, 250, 287, 293; 548/364.1, 364.7, 371.4, 372.5, 495; 549/68; 558/391, 392, 401, 402

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS ONLINE

search terms: structure searches

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP, A, 0,520,722, (BARKER ET AL.) 30 December 1992, see Table II, compound 1.	2, 8, 12
A	J. Med. Chem., Volume 26, issued 1983, Stout et al., "Synthesis and Antiarrhythmic and Parasympatholytic Properties of Substituted Phenols. 1. Heteroarylamine Derivatives", pages 808-813, see Table I, compound 2.	2, 8
X	Chemical Abstracts, Volume 117, issued 1990, Kokosi et al., "Process for producing imidazo[5,1-b]quinazolin-9-one derivatives and pharmaceutically acceptable salts thereof, as well as pharmaceutical compositions comprising such as active ingredient", abstract no. 212519, Hungarian patent 59411, see formula I.	3, 8, 10, 11

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance		
*E* earlier document published on or after the international filing date	X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*O* document referring to an oral disclosure, use, exhibition or other means		
*P* document published prior to the international filing date but later than the priority date claimed	&*	document member of the same patent family

Date of the actual completion of the international search 13 JUNE 1995	Date of mailing of the international search report 27 JUN 1995
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer Mary E. Ceperley <i>Mary E. Ceperley</i> Telephone No. (703) 308-0196

## INTERNATIONAL SEARCH REPORT

 International application No.  
 PCT/US95/01751

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Anti-Cancer Drugs, Volume 5, issued 1994, Kaur et al., "Tyrphostin induced growth inhibition: correlation with effect on p210 <sup>bcr-abl</sup> autokinase activity in K562 chronic myelogenous leukemia", pages 213-222, see Table 2, compound AG1112.	6, 8
A	JP, A, 55-167,205, 26 December 1980, see compounds 16 and 19.	8
A	DE, A, 1,135,471, (DEUSCHEL ET AL.) 30 August 1962, see claim 1.	8
X	Journal f. prakt. Chemie., Volume 329, No. 1, issued 1987, Vogel et al., "Synthese von 4-substituierten 1,2,3-Triazol[1,5-a]-chinoxalinen", pages 101-107, see compound 1f.	8
A	JP, A, 60-97964, 31 May 1985, see compound 1 and table of page 607.	8
X	Journal of Medicinal Chemistry, Volume 36, issued 1993, Gazit et al., "Tyrphostins. 3. Structure-activity relationship studies of $\alpha$ -substituted benzylidenemalononitrile 5-S-aryltyrphostins", pages 3556-3564, see Table II, compound 25.	8
X	Biochemistry, Volume 32, No. 17, issued 1993, Ohmichi et al., "The tyrosine kinase inhibitor tyrphostin blocks the cellular actions of nerve growth factor", pages 4650-4658, see page 4653, Table I, compounds Ag 18, 974, 1034, 1049.	8-11
X	Biochimica et Biophysica Acta, Volume 935, issued 1988, Miyoshi et al., "Quantitative analyses of the uncoupling activity of substituted phenols with mitochondria from flight muscles of house flies", pages 312-321, see Table I, compound 12.	8
X	Canadian Journal of Chemistry, Volume 63, issued 1985, Merlin, "Resonance Raman study of phenylhydrazonopropanedinitriles", pages 1840-1844, see abstract.	8

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/01751

## A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

A61K 31/24, 31/42, 31/275, 31/415, 31/495; C07C 211/45, 255/01; C07D 209/18, 231/38, 241/36, 265/34, 471/02;  
G01N 33/567

## A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

435/7.21; 436/503; 514/94, 249, 250, 287, 293; 548/364.1, 364.7, 371.4, 372.5, 495; 549/68; 558/391, 392, 401, 402